

Lab goals for Day 2

- Generalize your ODE code to handle systems of ODEs, solve the 2-body problem in GR for point particles in the “Newtonian orbits + quadrupole formula energy loss” approximation.
- Carry out a convergence test and evaluate the numerical error.
- How much energy is radiated in the inspiral?

post-Newtonian black holes

- Start with energy, e.g. as function of separation R or orbital frequency ω : $E(R)$, $E(\omega)$. Kepler: $\omega^2 R^3 = G M$.
- PN expansion:

$$\omega^2(R) = \frac{GM}{R^3} \left(1 + f_1(R) \left(\frac{v}{c}\right)^2 + f_2(R) \left(\frac{v}{c}\right)^4 + \dots \right)$$

- Compute energy loss $P = -dE/dt$ to some order in v/c , e.g. at leading order quadrupole formula (see GR text books like Wald)
- To compute the rate of change of any quantity X (e.g. $X = \omega, R$) we write

$$\frac{dX}{dt} = \frac{dE}{dX} \frac{dE}{dt}$$

- To lowest (Newtonian/quadrupole) order:

$$E(R) = m_1 + m_2 - M \frac{\eta}{2} \frac{M}{R}$$

$$E(\omega) = m_1 + m_2 - M \frac{\eta}{2} \left(\frac{(M\omega)^2}{G} \right)^{\frac{1}{3}}$$

$$\frac{dE}{dt} = -\frac{32}{5} \frac{G^4}{c^5} \eta^2 \left(\frac{v}{c} \right)^{10} (1 + O(v^2) + \dots)$$

- Here v is the velocity parameter, η the symmetric mass ratio:

$$v = (GM\omega)^{1/3}$$

$$\eta = \frac{m_1 m_2}{(m_1 + m_2)^2}$$

- For GW science, we also need the phase

$$\frac{d\phi}{dt} = \omega$$

exact solution

$$R(t) = \left(\frac{256}{5} \eta M^3 \right)^{\frac{1}{4}} (t_c - t)^{\frac{1}{4}}$$