

Leptoquarks and recent anomalies in the Flavour Sector



Rukmani Mohanta

School of Physics

University of Hyderabad

Hyderabad – 500046, INDIA


Based on: S. Sahoo and RM, PRD 91, 094019 (2015);
S. Sahoo and RM, PRD 93, 114001 (2016);
S. Sahoo, RM and A. Giri, PRD 95, 035027 (2017).

Outline

- Observed anomalies in the B sector
 - a. **$b \rightarrow s l^+ l^-$** (Neutral Current)
 - $B_s \rightarrow \mu^+ \mu^-$: Recent result and its Implications
 - $B \rightarrow K^* \mu^+ \mu^-$ measurements and anomalies
 - $B_s \rightarrow \phi \mu^+ \mu^-$ measurement
 - b. **$b \rightarrow c \tau \nu$** (Charged Current)
 - $B \rightarrow D^{(*)} \tau \nu$ processes
- LQ model as a possible NP candidate
- Conclusion

$b \rightarrow s l^+ l^-$ transitions


- Rare B decays mediated by FCNC $b \rightarrow s$ are sensitive probe for NP
- These decays occur via box or penguin diagrams in the SM (loop suppressed)
- CKM suppressed ($\propto V_{tb} V_{ts}^*$)
- New Physics can possibly be seen at the same level of SM
- Allow indirect search for new particles



Effective Hamiltonian for $b \rightarrow s$ transition


- The most general effective Hamiltonian describing $b \rightarrow s$ transition in OPE is

$$H_{eff} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_i \left[C_i(\mu) O_i(\mu) + C'_i(\mu) O'_i(\mu) \right] \rightarrow \text{RH suppressed in SM}$$



Electromagnetic penguin


$$Q_7^{(')} = \frac{e}{16\pi^2} m_b (\bar{s} \sigma_{\mu\nu} P_{R(L)} b) F^{\mu\nu}$$



Semi-leptonic vector current
Semi-leptonic A-V current

$$Q_9^{(')} = \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_{L(R)} b) (\bar{l} \gamma^\mu l)$$

$$Q_{10}^{(')} = \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_{L(R)} b) (\bar{l} \gamma^\mu \gamma_5 l)$$



Scalar


Pseudo-scalar

$$Q_S^{(')} = \frac{e^2}{16\pi^2} (\bar{s}_{L(R)} b_{R(L)}) (\bar{l} l)$$

$$Q_P^{(')} = \frac{e^2}{16\pi^2} (\bar{s}_{L(R)} b_{R(L)}) (\bar{l} \gamma_5 l)$$


*four quark operators $Q_{1\dots 6}$ only contribute through operator mixing.

Sensitivity to New Physics



$$\sim \frac{g^2}{16\pi^2} \frac{1}{M_W^2} V_{tb} V_{ts}^*$$

SM Amplitude is loop suppressed and CKM suppressed



$$\sim \frac{1}{\Lambda_{NP}^2}$$

Generic NP contributions not necessarily suppressed



$$\Lambda_{NP} \sim \frac{M_W}{g^2} \sqrt{\frac{16\pi^2}{|V_{tb} V_{ts}^*|}} \sim 10 \text{ TeV}$$

Rare B decays can probe high scales, potentially sensitive to NP, beyond the direct search at LHC.

I. Lepton Flavour Universality Violation (LFUV)


LHCb : $R_K = \text{BR}(B \rightarrow K \mu\mu) / \text{BR}(B \rightarrow K ee)$

in low q^2 region (1-6) GeV^2

$$R_K = 0.745^{+0.090}_{-0.074}(\text{stat}) \pm 0.036(\text{syst})$$

$$R_K = 1.0003 \pm 0.0001$$

There is 2.6 σ deficit from SM prediction



$$\text{Br}(B^+ \rightarrow K^+ \mu\mu)|_{[1-6]} = (1.19 \pm 0.07) \cdot 10^{-7}$$

$$\text{Br}(B^+ \rightarrow K^+ \mu\mu)|_{[1-6]}^{SM} = (1.75^{+0.60}_{-0.29}) \cdot 10^{-7}$$


- $\text{BR}(B^+ \rightarrow K^+ ee)$ in agreement with SM
- Can be explained if possible NP contributes to $b \rightarrow s \mu\mu$ not to $b \rightarrow s ee$.

II. $R_{K^*} = \text{BR}(B \rightarrow K^* \mu \mu) / \text{BR}(B \rightarrow K^* e e)$

- In April 2017, LHCb reported the measurement of R_{K^*} in two bins


$$R_{K^*}^{\text{LHCb}} = 0.660_{-0.070}^{+0.110} \pm 0.024, \quad q^2 \in [0.045, 1.1] \text{ GeV}^2 \quad \boxed{2.2\sigma}$$

$$R_{K^*}^{\text{LHCb}} = 0.685_{-0.069}^{+0.113} \pm 0.047, \quad q^2 \in [1.1, 6] \text{ GeV}^2 \quad \boxed{2.4\sigma}$$



III. $B_s \rightarrow \mu\mu$

- Highly suppressed in SM, (CKM & Helicity suppressed)
- Very clean experimental signature, involve only f_B



$$\begin{aligned}\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) &= (3.66 \pm 0.23) \times 10^{-9} \\ \mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) &= (1.06 \pm 0.09) \times 10^{-10}\end{aligned}$$

$$\begin{aligned}\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) &= (2.8^{+0.7}_{-0.6}) \times 10^{-9} \\ \mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) &= (3.9^{+1.6}_{-1.4}) \times 10^{-10}\end{aligned}$$





Precise SM predictions,
Bobeth et al, PRL 112,
101801 (2014)

Updated in 1702.05498

Combined LHCb and CMS
data

- Observation of $B_s \rightarrow \mu\mu$ at 6.2σ (Compatible with SM at 1.2σ)
- Observation of $B_d \rightarrow \mu\mu$, excess events at 3σ (Compatible with SM at 2.2σ)


III. $B_s \rightarrow \mu\mu$



Compatible with
SM at 2.3σ

$$\mathcal{R} \equiv B(B^0 \rightarrow \mu^+ \mu^-) / B(B_s^0 \rightarrow \mu^+ \mu^-).$$

LHCb Run I + Run II




$$\mathcal{B}(B_s \rightarrow \mu^+ \mu^-) = (3.0 \pm 0.6^{+0.3}_{-0.2}) \times 10^{-9}$$

$$\mathcal{B}(B_d \rightarrow \mu^+ \mu^-) < 3.4 \times 10^{-10}$$

$B_s \rightarrow \mu\mu$ (7.8σ) and $B_d \rightarrow \mu\mu$ (1.6σ)




ATLAS



$$\mathcal{BR}(B_s \rightarrow \mu^+ \mu^-) = 0.9^{+1.1}_{-0.8} \times 10^{-9} \quad (2\sigma)$$

$$\mathcal{BR}(B_d \rightarrow \mu^+ \mu^-) < 4.2 \times 10^{-10} \quad (95\%CL)$$

IV. Results on $b \rightarrow s \mu\mu$ branching fractions



Data is average of LHCb, CMS, ATLAS and CDF.


$$C_9 = C_9^{\text{SM}} - 1.0$$

$$C_9' = 1.2$$

Branching ratio tend to lie below SM predictions
Horgan et al., PRL 112, 212003 (2014)

V. Results from $B_d \rightarrow K^* \mu^+ \mu^-$

- The decay $B \rightarrow K^*(\rightarrow K\pi) \mu^+ \mu^-$ plays a special role
- Ang dist. of the 4-body final state gives numerous observables sensitive to NP.




- Forbidden at tree level
- Penguin and Box diagrams

- Assortment of measurables whose dependence on $\mu^+ \mu^-$ mass can reveal NP
- Angular distributions: Forward-Backward asymmetry and its zero point

- The decay distribution of B^0

$$\frac{d^4\Gamma}{dq^2 d\cos\theta_\ell d\cos\theta_K d\phi} = \frac{9}{32\pi} \left[I_1^s \sin^2\theta_K + I_1^c \cos^2\theta_K \right. \\ + I_2^s \sin^2\theta_K \cos 2\theta_\ell + I_2^c \cos^2\theta_K \cos 2\theta_\ell \\ + I_3 \sin^2\theta_K \sin^2\theta_\ell \cos 2\phi + I_4 \sin 2\theta_K \sin 2\theta_\ell \cos \phi \\ + I_5 \sin 2\theta_K \sin \theta_\ell \cos \phi + I_6 \sin^2\theta_K \cos \theta_\ell \\ + I_7 \sin 2\theta_K \sin \theta_\ell \sin \phi + I_8 \sin 2\theta_K \sin 2\theta_\ell \sin \phi \\ \left. + I_9 \sin^2\theta_K \sin^2\theta_\ell \sin 2\phi \right],$$



- Combining B^0 and $\bar{B}^0$ decays,

CP Avg.

$$S_j = (I_j + \bar{I}_j) \bigg/ \frac{d\Gamma}{dq^2} \quad \text{or} \quad A_j = (I_j - \bar{I}_j) \bigg/ \frac{d\Gamma}{dq^2}.$$

CP Asymmetry


S_7, S_8 and S_9 depend on combinations $\text{Im}(\mathcal{A}_m \mathcal{A}_n^*)$

- Suppressed by small size of strong phase and are expected to be zero for the full q^2 (A_7, A_8 and A_9 are not)

Result on P_5'

Less form-factor dependent observable:
(smaller theoretical uncertainties)



$$P_5' = \frac{S_5}{\sqrt{F_L(1 - F_L)}}$$



[Descotes-Genon et al, [JHEP 05 \(2013\) 137](#)]

- Two bins each show local deviations of $\sim 3\sigma$
(LHCb: 3 fb^{-1} data, 3.4 sigma deviation 1512.04442)
New Belle result consistent with LHCb

Belle Results on $P'_{4,5}$





Includes $B^0 \rightarrow K^{*0} \mu^+ \mu^-$, $B^+ \rightarrow K^{*+} \mu^+ \mu^-$, $B^0 \rightarrow K^{*0} e^+ e^-$, and $B^+ \rightarrow K^{*+} e^+ e^-$

1612.05014

$Q_{4,5}$ Observables:

$$Q_i = P_i^{\mu} - P_i^e$$




NP Scenario: $C_{9\mu}^{\text{NP}} = -1.1$ (1612.15014)

$b \rightarrow s \mu\mu$ Anomalies

Decay	obs.	q^2 bin	SM pred.	measurement		pull
$B^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	F_L	[2, 4.3]	0.81 ± 0.02	0.26 ± 0.19	ATLAS	+2.9
$B^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	F_L	[4, 6]	0.74 ± 0.04	0.61 ± 0.06	LHCb	+1.9
$\bar{B}^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	S_5	[4, 6]	-0.33 ± 0.03	-0.15 ± 0.08	LHCb	-2.2
$B^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	P'_5	[1.1, 6]	-0.44 ± 0.08	-0.05 ± 0.11	LHCb	-2.9
$\bar{B}^0 \rightarrow \bar{K}^{*0} \mu^+ \mu^-$	P'_5	[4, 6]	-0.77 ± 0.06	-0.30 ± 0.16	LHCb	-2.8
$B^- \rightarrow K^{*-} \mu^+ \mu^-$	$10^7 \frac{d\text{BR}}{dq^2}$	[4, 6]	0.54 ± 0.08	0.26 ± 0.10	LHCb	+2.1
$\bar{B}^0 \rightarrow \bar{K}^0 \mu^+ \mu^-$	$10^8 \frac{d\text{BR}}{dq^2}$	[0.1, 2]	2.71 ± 0.50	1.26 ± 0.56	LHCb	+1.9
$B^0 \rightarrow \bar{K}^0 \mu^+ \mu^-$	$10^8 \frac{d\text{BR}}{dq^2}$	[16, 23]	0.93 ± 0.12	0.37 ± 0.22	CDF	+2.2
$B_s \rightarrow \phi \mu^+ \mu^-$	$10^7 \frac{d\text{BR}}{dq^2}$	[1, 6]	0.48 ± 0.06	0.23 ± 0.05	LHCb	+3.1

arXiv:1503.06199

Best fit Results (arXiv: 1704.05340)



Global fit: $C_{9\mu}^{NP} = -1.1$


Other coefficients play a secondary role: $C_9^{NP}, C_9^{NP} = -C_{10}^{NP}, C_9^{NP} = -C_9^{NP}$

R_{D^*} Anomaly in $B \rightarrow D^{(*)} l \bar{\nu}$ (Tree level)

- Belle/BaBar/LHCb: $R_{D^{(*)}} = \frac{\Gamma(B \rightarrow D^{(*)} \tau \bar{\nu})}{\Gamma(B \rightarrow D^{(*)} l \bar{\nu})}$

$$R_D = 0.397 \pm 0.049 \text{ (World Ave.)}$$
$$= 0.30 \pm 0.01 \text{ (SM)}$$

$$R_{D^*} = 0.316 \pm 0.016 \text{ (World Ave.)}$$
$$= 0.252 \pm 0.005 \text{ (SM)}$$



$\sim 4\sigma$ deviation from SM predictions


NP competing with SM at tree level

Effective Hamiltonian for $b \rightarrow c \tau \nu$

$$\mathcal{H}_{\text{eff}} = \frac{4G_F}{\sqrt{2}} V_{cb} \left[\left(\delta_{l\tau} + C_{V_1}^l \right) \mathcal{O}_{V_1}^l + C_{V_2}^l \mathcal{O}_{V_2}^l + C_{S_1}^l \mathcal{O}_{S_1}^l + C_{S_2}^l \mathcal{O}_{S_2}^l \right],$$

where the operators are

$$\begin{aligned} \mathcal{O}_{V_1}^l &= (\bar{c}_L \gamma^\mu b_L) (\bar{\tau}_L \gamma_\mu \nu_L), & \mathcal{O}_{V_2}^l &= (\bar{c}_R \gamma^\mu b_R) (\bar{\tau}_L \gamma_\mu \nu_L), \\ \mathcal{O}_{S_1}^l &= (\bar{c}_L b_R) (\bar{\tau}_R \nu_L), & \mathcal{O}_{S_2}^l &= (\bar{c}_R b_L) (\bar{\tau}_R \nu_L). \end{aligned}$$



Leptoquarks

- LQs are color triplet bosons
- Coupled to Quarks and Lepton simultaneously
- Classified by $F = 3B + L$, $F = 0, 2$
- Spin=0,1
- Naturally emerge in many extensions of SM
- $|Q|=1/3, 2/3, 4/3, 5/3$
- Representation under SM Gauge group:
 - Triplet under SU(3): as Q(triplet) and L(singlet)
 - Singlet, Doublet, Triplet under SU(2): As LH quarks and leptons are doublets and RH are singlets
 - Y is characterized by the charge $Q = T_3 + Y$
- Intergenerational mixing is severely restricted by FCNC data


List of Scalar and Vector LQs (1603.04993)

$(SU(3), SU(2), U(1))$	Spin	Symbol	Type	F
$(\bar{\mathbf{3}}, \mathbf{3}, 1/3)$	0	S_3	$LL(S_1^L)$	-2
$(\mathbf{3}, \mathbf{2}, 7/6)$	0	R_2	$RL(S_{1/2}^L), LR(S_{1/2}^R)$	0
$(\mathbf{3}, \mathbf{2}, 1/6)$	0	$\tilde{R}_2$	$RL(\tilde{S}_{1/2}^L), \overline{LR}(\tilde{S}_{1/2}^L)$	0
$(\bar{\mathbf{3}}, \mathbf{1}, 4/3)$	0	$\tilde{S}_1$	$RR(\tilde{S}_0^R)$	-2
$(\bar{\mathbf{3}}, \mathbf{1}, 1/3)$	0	S_1	$LL(S_0^L), RR(S_0^R), \overline{RR}(S_0^R)$	-2
$(\bar{\mathbf{3}}, \mathbf{1}, -2/3)$	0	$\bar{S}_1$	$\overline{RR}(\bar{S}_0^R)$	-2
$(\mathbf{3}, \mathbf{3}, 2/3)$	1	U_3	$LL(V_1^L)$	0
$(\bar{\mathbf{3}}, \mathbf{2}, 5/6)$	1	V_2	$RL(V_{1/2}^L), LR(V_{1/2}^R)$	-2
$(\bar{\mathbf{3}}, \mathbf{2}, -1/6)$	1	$\tilde{V}_2$	$RL(\tilde{V}_{1/2}^L), \overline{LR}(\tilde{V}_{1/2}^R)$	-2
$(\mathbf{3}, \mathbf{1}, 5/3)$	1	$\tilde{U}_1$	$RR(\tilde{V}_0^R)$	0
$(\mathbf{3}, \mathbf{1}, 2/3)$	1	U_1	$LL(V_0^L), RR(V_0^R), \overline{RR}(V_0^R)$	0
$(\mathbf{3}, \mathbf{1}, -1/3)$	1	$\bar{U}_1$	$\overline{RR}(\bar{V}_0^R)$	0

Leptoquark Model

Wise et al, PRD 88,035009 (2013)

- $|F|$ can be either 0 or 2 depending on whether LQ couples to fermion-antifermion pair or fermion-fermion pair
- $F = 0$ interaction between leptoquark and quark-lepton pair due to a scalar doublet: $X = (3, 2, 7/6)$ and $X = (3, 2, 1/6)$ or vector $U^3_\mu (3, 3, 2/3)$ or singlets $U^1_\mu (3, 1, 2/3)$.
- Couplings are not lepton flavour universal or conserving



Scalar LQs and $b \rightarrow s$ processes

- Models with scalar leptoquarks can modify the $\mathbf{H}_{\text{eff}}$
- Minimal renormalizable scalar LQ model with **B and L conserving couplings** which do not allow proton decay
- $X = (3, 2, 7/6)$ and $X = (3, 2, 1/6)$ under the $SU(3) \times SU(2) \times U(1)$
- Interaction Lagrangian for the scalar leptoquark $X = (3, 2, 7/6)$

$$\mathcal{L} = -\lambda_u^{ij} \bar{u}_R^i X^T \epsilon L_L^j - \lambda_e^{ij} \bar{e}_R^i X^\dagger Q_L^j + h.c.,$$

$$X = \begin{pmatrix} V_\alpha \\ Y_\alpha \end{pmatrix}, \quad \epsilon = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

- After expanding the $SU(2)$ indices and Fierz transformation,

$$\mathcal{H}_{LQ} = \frac{\lambda_\mu^{32} \lambda_\mu^{22*}}{8M_Y^2} [\bar{s} \gamma^\mu (1 - \gamma_5) b] [\bar{\mu} \gamma_\mu (1 + \gamma_5) \mu] = \frac{\lambda_\mu^{32} \lambda_\mu^{22*}}{4M_Y^2} (\mathcal{O}_9 + \mathcal{O}_{10}),$$

- One obtains the new Wilson coefficients


$$C_9^{NP} = C_{10}^{NP} = -\frac{\pi}{2\sqrt{2}G_f\alpha V_{tb}V_{ts}^*} \frac{\lambda_\mu^{32}\lambda_\mu^{22*}}{M_Y^2}.$$

- Lagrangian for the coupling of scalar leptoquark $X = (3, 2, 1/6)$

$$\mathcal{L} = -\lambda_d^{ij} \bar{d}_R^i X^T \epsilon L_L^j + h.c.,$$

- Which after expansion gives



$$\mathcal{H}_{LQ} = \frac{\lambda_s^{22}\lambda_b^{32*}}{8M_V^2} [\bar{s}\gamma^\mu(1+\gamma_5)b] [\bar{\mu}\gamma_\mu(1-\gamma_5)\mu] = \frac{\lambda_s^{22}\lambda_b^{32*}}{4M_V^2} (\mathcal{O}'_9 - \mathcal{O}'_{10}),$$



$$C_9'^{NP} = -C_{10}'^{NP} = \frac{\pi}{2\sqrt{2}G_f\alpha V_{tb}V_{ts}^*} \frac{\lambda_s^{22}\lambda_b^{32*}}{M_V^2}.$$

LQ Parameter space: $B_s \rightarrow l^+ l^- B_s$, B_s - B_s mixing $B \rightarrow X_s l^+ l^-$ both at high q^2 and low q^2

P_4' and P_5'



S. Sahoo & RM, PRD 93, 034018 (2016)

This simple model can explain most of the anomalies

Lepton Flavour Universality

Recently LHCb measured the ratio of decay rates $B \rightarrow K \mu \mu$ and $B \rightarrow K e e$ in low q^2 region

$$R_K = 0.745^{+0.090}_{-0.074}(\text{stat}) \pm 0.036(\text{syst})$$



This ratio is expected to be 1 in the SM with uncertainty less than 1%.

There is 2.6σ deficit from SM prediction



- This result can be accommodated in the scalar LQ model with predicted value

$$R_K = 0.62 - 0.96$$

S Sahoo and R. Mohanta, PRD 094019 (2015)



$B \rightarrow K \mu\mu$ in high q^2



Variation of differential BR and R_K in high q^2 region

S. Sahoo and R Mohanta, NJP 18, 013032 (2016)

Explanation to R_K and $R_{D^{(*)}}$ with scalar LQ

- $(3,2,1/6)$ can explain both these anomalies for RH neutrinos (two states with $Q = 2/3, -1/3$)

$$\mathcal{L}_Y = -x_{ij}\bar{d}_R^i e_L^j \tilde{R}_2^{2/3} + (xV_{\text{PMNS}})_{ij}\bar{d}_R^i \nu_L^j \tilde{R}_2^{-1/3} \\ + (V_{\text{CKM}}y)_{ij}\bar{u}_L^i \nu_R^j \tilde{R}_2^{2/3} + y_{ij}\bar{d}_L^i \nu_R^j \tilde{R}_2^{-1/3} + \text{h.c.}$$

$$R_K: C_9' = -C_{10}'$$


- Since LQ couples to RH neutrinos there will be no interference with SM

$$|\mathcal{M}(B \rightarrow D^{(*)}\ell\nu)|^2 = |\mathcal{M}_{\text{SM}}|^2 + |\mathcal{M}_{\text{NP}}|^2$$

Combination of two scalar LQs

- One can consider two LQs with same mass (one singlet and other triplet under SU(2))

$$L = \lambda_{fi}^{1L} \overline{Q}_f^c i\tau_2 L_i \Phi_1^\dagger + \lambda_{fi}^{3L} \overline{Q}_f^c i\tau_2 (\tau \cdot \Phi_3)^\dagger L_i + \text{h.c.}.$$



Vector LQ and B anomalies

- We consider the LQs $U_1(3,1,2/3)$, $U_3(3,3,2/3)$, which can mediate $b \rightarrow c \ell \nu$ and $b \rightarrow s \ell \ell$
- Conserve baryon number, avoid rapid proton decay
- $\mathcal{L}_{\text{int}}$ of $U_{1,3}$ LQs with SM fermion bilinear is

$$\mathcal{L}^{LQ} = (h_{1L}^{ij} \bar{Q}_{iL} \gamma^\mu L_{jL} + h_{1R}^{ij} \bar{d}_{iR} \gamma^\mu l_{jR}) U_{1\mu} + h_{3L}^{ij} \bar{Q}_{iL} \boldsymbol{\sigma} \gamma^\mu L_{jL} \mathbf{U}_{3\mu}$$

- Int. Lagrangian gives additional contributions to $b \rightarrow c \tau \nu$:

$$C_{V_1}^{LQ} = \frac{1}{2\sqrt{2}G_F V_{cb}} \sum_{k=1}^3 V_{k3} \left[\frac{h_{1L}^{2l} h_{1L}^{k3*}}{M_{U_1^{2/3}}^2} - \frac{h_{3L}^{2l} h_{3L}^{k3*}}{M_{U_3^{2/3}}^2} \right],$$

$$C_{V_2}^{LQ} = 0, \quad C_{S_1}^{LQ} = -\frac{1}{2\sqrt{2}G_F V_{cb}} \sum_{k=1}^3 V_{k3} \frac{2h_{1L}^{2l} h_{1R}^{k3*}}{M_{U_1^{2/3}}^2}.$$

- It also gives additional contributions to $b \rightarrow s \ell \ell$

$$C_9^{\text{NP}} = -C_{10}^{\text{NP}} = \frac{\pi}{\sqrt{2}G_F V_{tb} V_{ts}^* \alpha} \left[\frac{h_{1L}^{2l} h_{1L}^{k3*}}{M_{U_1^{2/3}}^2} + \frac{h_{3L}^{2l} h_{3L}^{k3*}}{M_{U_3^{2/3}}^2} \right],$$

$$C_9'^{\text{NP}} = C_{10}'^{\text{NP}} = \frac{\pi}{\sqrt{2}G_F V_{tb} V_{ts}^* \alpha} \frac{h_{1R}^{2l} h_{1R}^{k3*}}{M_{U_1^{2/3}}^2},$$

$$-C_P^{\text{NP}} = C_S^{\text{NP}} = \frac{\sqrt{2}\pi}{G_F V_{tb} V_{ts}^* \alpha} \frac{h_{1L}^{2l} h_{1R}^{k3*}}{M_{U_1^{2/3}}^2},$$

$$C_P'^{\text{NP}} = C_S'^{\text{NP}} = \frac{\sqrt{2}\pi}{G_F V_{tb} V_{ts}^* \alpha} \frac{h_{1R}^{2l} h_{1L}^{k3*}}{M_{U_1^{2/3}}^2},$$

LQ parameter space
constrained from:



$B_s \rightarrow l^+ l^-$

$B \rightarrow X_s \nu \nu$

$B \rightarrow X_s l^+ l^-$ both at high q^2
and low q^2

Muon (g-2)



Results: $R_{D(*)}$



Observables	SM Predictions	Values in LQ Model	Experimental Limit
R_D	0.31	0.11 – 0.386	$0.397 \pm 0.040 \pm 0.028$
R_{D^*}	0.26	0.243 – 0.32	$0.316 \pm 0.016 \pm 0.010$

S. Sahoo, R. Mohanta and A. K. Giri, PRD, 95, 035027 (2017).



Results: R_K



Observables	SM Predictions	Values in LQ Model	Experimental Limit
$R_K^{\mu e}_{q^2 \in [1, 6]}$	1.006	0.75 – 1.006	$0.745^{+0.090}_{-0.074} \pm 0.036$
$R_K^{\mu e}_{q^2 \geq 14.18}$	1.004	0.74 – 1.004	...



S. Sahoo, R. Mohanta and A. K. Giri, PRD, 95, 035027 (2017).

R_{K^*}





Observables	SM Predictions	Values in LQ Model
$R_{K^*}^{\mu e}$ $q^2 \in [0.045, 1.1]$	0.996	0.725 – 0.996
$R_{K^*}^{\mu e}$ $q^2 \in [1.1, 6]$	0.996	0.725 – 0.996
$R_{K^*}^{\mu e}$ $q^2 \geq 14.18$	0.999	0.816 – 0.999

P_5'



$Q_{4,5}$



Summary

- There exist some mild discrepancies in the rare decays involving $b \rightarrow s$ processes.
- Intriguing hints for Lepton Flavour Universality violating New Physics
- It is not yet confirmed whether these are the smoking gun signals of some NP or just the statistical fluctuations.
- Scalar/Vector LQ models can accommodate these anomalies.
- Hopefully future results from the Flavour Factories will settle these issues and/or provide the new direction to High Energy Physics.

THANK YOU