

Moduli Masses and Inflationary Observables

Koushik Dutta
Theory Division, Saha Institute

Candles of Darkness, ICTS, Bangalore

June, 2017

References

Moduli Vacuum Misalignment and Precise Predictions in String Inflation

[arXiv: 1604.08512\[hep-th\]](#)

Michele Cicoli, Koushik Dutta, Anshuman Maharana, Fernando Quevedo

Inflationary Predictions and Moduli Masses

[arXiv:1506.05745\[hep-ph\]](#)

Kumar Das, Koushik Dutta, Anshuman Maharana

Inflationary constraints on modulus dominated cosmology

[arXiv:1409.7037\[hep-ph\]](#)


Koushik Dutta, Anshuman Maharana

Take Home

The number of e-foldings is sensitive to the post-inflationary history of the universe.

The generic presence of light scalar fields (in SUSY/String Theory) leads to a late-time period of matter domination which lowers the required number of e-foldings and, in turn, modifies the exact predictions of any inflationary model.


CMB



$$\frac{\Delta T(\hat{n})}{\bar{T}} = \sum_{l=0}^{\infty} \sum_{m=-l}^{m=+l} a_{lm} Y_{lm}(\hat{n})$$


$$C(\theta) \equiv \left\langle \frac{\Delta T}{\bar{T}}(\hat{n}) \frac{\Delta T}{\bar{T}}(\hat{n}') \right\rangle$$

$$C_l^{obs} = \frac{1}{2l+1} \sum_m |a_{lm}|^2$$






CMB Data

Photons are linearly polarised



$$\langle a_{X,lm} a_{Y,l'm'}^* \rangle = C_l^{XY} \delta_{ll'} \delta_{mm'}, \quad X, Y = T, E, B$$



$$\mathcal{D}_l = l(l+1)C_l/2\pi$$

CMB Theory

Strongly coupled plasma - sound waves

(red line in previous plot) $C_l = \int d \ln k \Delta_{\mathcal{R}}^2(k) T_l^2(k)$

primordial spectrum

A_s, n_s (given in radiation dominated era)

transfer function

$\Omega_b h^2, \Omega_c h^2, \Omega_\Lambda, \tau$

$$\Delta_{\mathcal{R}}^2(k) = A_s (k/k_*)^{n_s-1}$$

Λ CDM cosmology



Parameter	Planck	... + WMAP + ACT	CMB + BAO
$\Omega_b h^2$	0.02207 ± 0.00067	0.02207 ± 0.00054	0.02214 ± 0.00048
$\Omega_c h^2$	0.1196 ± 0.0061	0.1198 ± 0.0052	0.1197 ± 0.0034
Ω_Λ	0.683 ± 0.040	0.685 ± 0.033	0.692 ± 0.021
τ	0.097 ± 0.080	0.091 ± 0.027	0.092 ± 0.026
$10^9 A_s$	2.23 ± 0.32	2.20 ± 0.11	2.20 ± 0.11
n_s	0.962 ± 0.019	0.959 ± 0.014	0.961 ± 0.011

Complete story

Coherent Phase

(of sub-Hubble modes)

given an wavenumber, infinite number of modes





with random phase

This is **still** sub-horizon physics

Coherent Phase

(of super-Hubble modes)



$$D_l^{TE} \sim \langle Q_{lm}^T Q_{lm}^E \rangle$$

$$\Delta T \sim \delta_\gamma, \quad E \sim \partial v_\gamma$$

$$\delta_\gamma = A \cos(\omega t + \phi) e^{i\vec{k} \cdot \vec{x}}$$

$$\partial v_\gamma = -\dot{\delta}_\gamma$$

If phases were
uniformly distributed

$$D_l^{TE} = 0$$



phase is not
random

From data ONLY

- At the time of BBN, the Universe was radiation dominated
- The existence of primordial spectrum

$$\Delta_{\mathcal{R}}^2(k) = A_s (k/k_*)^{n_s-1}$$

- Coherent super-Hubble perturbations
- (possibly) dark matter .. gravitational collapse

From data ONLY

- At the time of BBN, the Universe was radiation dominated
- The existence of primordial spectrum

$$\Delta_{\mathcal{R}}^2(k) = A_s (k/k_*)^{n_s-1}$$

- Coherent super-Hubble perturbations
- (possibly) dark matter .. gravitational collapse


INFLATION

INFLATION also solves horizon and flatness problem

From: Kane, Sinha, Watson (2015)

Thermal History

Alternative History



Outline


- Cosmology with moduli
- Constraint on moduli mass/inflation
Dutta, Maharana (2014)
- Implications ..
Das, Dutta, Maharana (2015)
- An example: Kahler moduli inflation
Cicoli, Dutta, Maharana, Quevedo (2016)

Framework

'Modular Cosmology'

Moduli φ

potential at
the end of
inflation



$$\varphi = 0$$

V minima during
inflation


$$\hat{\varphi} \sim M_{Pl}$$

post-inflationary


moduli mass $m < H_{inf}$

$$Y = \hat{\varphi}/M_{Pl} \sim 1$$

Inflation Observables




Inflation Observables



Few observables and other observables
seem to be not required

Where we stand





$$V^{1/4} \sim \left(\frac{r}{0.01} \right)^{1/4} 10^{16} \text{ GeV}$$

We need some luck here!


Future

CORE/LITEBIRD



precise measurements of spectral index is crucial! sensitivity $n_s \sim 10^{-3}$

Thermal History



Consistency

V_k must be evolved to H_0

Any post inflationary phase must be evolved to the present energy density

Making predictions ..

$$N_{inf} + \frac{1}{4}(1 - 3w_{rh})N_{rh} = 55 + \frac{1}{4}\ln r + \frac{1}{4}\ln(\rho_k/\rho_{end})$$

$$N_{inf} = 55 \pm 5$$

Liddle, Leach (2003)

PLANCK (2016)


'Theoretical prior'

compute observables in terms of N_{inf} and
see whether it fits data for $N = 50-60$!

$$V(\chi) = \frac{1}{2}m^2\chi^2$$

$$n_s - 1 = -\frac{2}{N_k}$$

$$r = 8/N_K$$



'Theoretical prior'

How does making
predictions change for
modular cosmology?

Moduli ..

- moduli: light scalar fields with Planck suppressed interactions
- at tree level effective Lagrangian of string theory/SUGRA, moduli are massless
- moduli must acquire masses (thus fixed vev) to become phenomenologically viable
- moduli stabilisation: KKLT etc....

Moduli ..

- Conservative approach: Make ALL modulus much heavier than the Hubble scale ..
decouple from inflation
- Wishful
- In practice, few fields remain parametrically light in the post-inflationary vacua .. (e.g LVS constructions .. / talk by John Ellis yesterday) see later ..


A typical case

$$\mathcal{L} \supset -\frac{1}{2}m^2\varphi^2 - \frac{1}{2}H^2(\varphi - \hat{\varphi})^2 - V_{inf}(\chi)$$

$$m < H_{inf}$$

post-inflationary
moduli mass

minima during
inflation



$$Y = \hat{\varphi}/M_{Pl} \sim 1$$

Dine, Randall, Thomas

Supergravity

$$V = e^{K[\varphi, \bar{\varphi}]} V_0[\varphi, \chi] \sim H^2 M_{Pl}^2 f \left(\frac{\varphi}{M_{Pl}} \right)$$

$$V'' \sim H^2$$

η – problem

Scale of variations M_{Pl}

$$Y = \hat{\varphi}/M_{Pl} \sim 1$$

Dine, Randall, Thomas

Dvali

Supergravity

$$V = e^{K[\varphi, \bar{\varphi}]} V_0[\varphi, \chi] \sim H^2 M_{Pl}^2 f \left(\frac{\varphi}{M_{Pl}} \right)$$

$$V'' \sim H^2$$

η – problem

Scale of variations M_{Pl}

$$Y = \hat{\varphi}/M_{Pl} \sim 1$$

Dine, Randall, Thomas

Toy example ..

Dvali

$$V = (m_{3/2}^2 - a^2 H^2) |\varphi|^2 + \frac{1}{2M_{Pl}^2} (m_{3/2}^2 + b^2 H^2) |\varphi|^4$$

$$\hat{\varphi} \sim (a/b) M_{Pl}$$

Different source of moduli stabilisation
during and after inflation

Antusch, K.D, Haller

Decay of Modulus

moduli must decay so that it does not overclose the Universe

$$\Gamma_{mod} \sim \frac{m_\phi^3}{16\pi M_{Pl}^2}$$

G. D. Coughlan, W. Fischler,
E. W. Kolb, S. Raby and G. G.
Ross 1984

$$\rho_{mod}(t_{decay}) = g \frac{\pi^2}{30} T_{reheat}^4 = 3H^2 M_{Pl}^2$$

↑
 Γ_{mod} decay happens

$$T_{reheat} \sim \sqrt{\Gamma M_{Pl}} \quad T_{reheat} > MeV \quad \text{successful BBN}$$

$$m_\phi > 30 \text{ TeV} \quad \text{BBN bound}$$

phenomenological implications .. SUSY
breaking ..

Inflation $\longrightarrow$ reheating $\longrightarrow$ radiation

N_{inf} N_{rh1} N_{rad}

reheating $\longleftarrow$ moduli (matter)

N_{rh2} 'non-standard' N_{mod}

BBN

today

Inflation $\longrightarrow$ reheating $\longrightarrow$ radiation

N_{inf} N_{rh1} N_{rad}

reheating $\longleftarrow$ moduli (matter)

N_{rh2} 'non-standard' N_{mod}

BBN

today

K.D, Maharana

arXiv:1409.7037[hep-ph]

Liddle, Leach (2003)

$$N_{inf} + \frac{1}{4}(1 - 3w_{rh1})N_{rh1} + \frac{1}{4}N_{mod} + \frac{1}{4}(1 - 3w_{rh2})N_{rh2}$$

$$= 55.43 + \frac{1}{4}\ln r + \frac{1}{4}\ln\left(\frac{\rho_k}{\rho_{end}}\right)$$

non-thermal history

Constraint ..

$$\Gamma_{mod} \sim \frac{m_\varphi^3}{16\pi M_{Pl}^2}$$

$$N_{mod} \sim \frac{2}{3} \ln\left(\frac{16\pi M_{Pl}^2 Y^4}{m_\phi^2}\right)$$

initial displacement $Y = \hat{\varphi}/M_{Pl}$

$$\frac{1}{6} \ln\left(\frac{16\pi M_{Pl}^2 Y^4}{m_\varphi^2}\right) + \frac{1}{4}(1 - 3w_{rh1})N_{re1} + \frac{1}{4}(1 - 3w_{rh2})N_{re2}$$

post-inflationary details

$$= 55.43 - N_{inf} + \frac{1}{4} \ln r + \frac{1}{4} \ln\left(\frac{\rho_k}{\rho_{end}}\right)$$

inflationary details

inflationary
potentials

Implications I

Central value of e-folding shifts

$$N_{inf} = 55 \pm 5$$



$$N_{inf} = \left(55 - \frac{1}{3} \ln \left(\frac{\sqrt{16\pi} M_{pl} Y^2}{m_\varphi} \right) \right) \pm 5$$

For $m_\varphi \sim 10^3$ TeV :


$$N_{inf} = 41 \quad - \quad 51$$



For $m_\varphi \sim 10^6$ TeV :

$$N_{inf} = 43 \quad - \quad 53$$

(used to be 50 - 60)

($Y \sim 0.1$ assumed)





Das, K.D, Maharana

sensitivity $n_s \sim 10^{-3}$

CMB IV, CORE

Implications

- moduli masses tied to soft masses in SUSY
- typical moduli mass 100/1000 TeV

Preferred value of inflation e-folds

$$\hat{N}_{inf} = 55 - \frac{1}{3} \left(\frac{\sqrt{16\pi} M_{Pl} Y^2}{m_\varphi} \right) \simeq 45$$

The effects of modulus mass must be taken for inflation models for $m_\varphi \lesssim 10^{10} \text{ GeV}$

Implications II

$$\frac{1}{6} \ln\left(\frac{16\pi M_{Pl}^2 Y^4}{m_\varphi^2}\right) + \frac{1}{4}(1 - 3w_{rh1})N_{re1} + \frac{1}{4}(1 - 3w_{rh2})N_{re2}$$

usually positive definite

$$= 55.43 - N_{inf} + \frac{1}{4} \ln r + \frac{1}{4} \ln\left(\frac{\rho_k}{\rho_{end}}\right)$$

analytical/numerical
understanding of
reheating: $w_{re} < 1/3$

Ellis, Garcia, Nanopoulos, Olive (2015)

Constraint ..

$$m_\varphi \gtrsim \sqrt{16\pi} M_{\text{pl}} Y^2 e^{-3\left(55.43 - N_k + \frac{1}{4} \ln\left(\frac{\rho_k}{\rho_{\text{end}}}\right) + \frac{1}{4} \ln r\right)}$$

- Dependence correlated

Das, K.D, Maharana

- Larger the value of $N_{\{k\}}$, stronger the bound

- smaller the value of 'r' stronger the bound

- bound depends on the nature of inflationary potentials via the ratio of energy densities

Small field models


$$m_\varphi \gtrsim \sqrt{16\pi} M_{\text{pl}} Y^2 e^{-3\left(55.43 - N_k + \frac{1}{4} \ln\left(\frac{\rho_k}{\rho_{\text{end}}}\right) + \frac{1}{4} \ln r\right)}$$

- conservative estimate $r \sim 0.01$ stronger the bound
- potential plateau like .. ratio of energy densities negligible
- take $Y = 0.1$, then for $N = 50$

$$m_\varphi \gtrsim 4.5 \times 10^6 \text{ TeV}$$

much stronger than BBN bound

Small field models



for $N > 44$,
the bound
much stronger
than BBN
bound

Das, Maharana, K.D

large field models

$$m_\varphi \gtrsim \sqrt{16\pi} M_{\text{pl}} Y^2 e^{-3\left(55.43 - N_k + \frac{1}{4} \ln\left(\frac{\rho_k}{\rho_{\text{end}}}\right) + \frac{1}{4} \ln r\right)}$$

$$V_\chi = m^{4-\alpha} \chi^\alpha$$

chaotic inflation
axion monodromy

$$m_\phi \gtrsim$$

$$\sqrt{16\pi} M_{\text{pl}} Y^2 e^{-3\left(55.85 - \frac{(2+\alpha)}{2(1-n_s)} + \frac{\alpha}{8} \ln 2 + \frac{1}{8} (\alpha-2) \ln\left(\frac{2+\alpha}{\alpha(1-n_s)}\right)\right)}$$

$$\alpha = 2 \quad m_\varphi \gg 10^{10} \text{ TeV} \quad \text{PLANCK: Central value}$$

$$\alpha = 2/3$$

Bound insignificant

What to calculate now?

Cicoli, K.D, Maharana, Quevedo


$$N_{mod} \sim \frac{2}{3} \ln\left(\frac{16\pi M_{Pl}^2 Y^4}{m_\phi^2}\right)$$

$$\begin{aligned} & \frac{1}{6} \ln\left(\frac{16\pi M_{Pl}^2 Y^4}{m_\phi^2}\right) + \frac{1}{4}(1 - 3w_{rh1})N_{re1} + \frac{1}{4}(1 - 3w_{rh2})N_{re2} \\ &= 55.43 - N_{inf} + \frac{1}{4} \ln r + \frac{1}{4} \ln\left(\frac{\rho_k}{\rho_{end}}\right) \end{aligned}$$

Kahler moduli inflation

Conlon, Quevedo (2005)

- In LVS scenario of string theory.
- A concrete set-up where inflationary potential is known
- Post-inflationary modulus domination happens



$$\Gamma_{\tau_n} \simeq 0.1 \frac{m_{\tau_n}^3}{M_s^2} \gg \Gamma_\nu \simeq \frac{m_\nu^3}{16\pi M_{pl}^2}$$

Cicoli, Mazumdar (2010)

Details ..

Volume $\mathcal{V} = \alpha \left(\tau_1^{3/2} - \sum_{i=2}^n \lambda_i \tau_i^{3/2} \right)$

$$V = \sum_{i=2}^n \frac{8(a_i A_i)^2 \sqrt{\tau_i}}{3\mathcal{V} \lambda_i} e^{-2a_i \tau_i} - \sum_{i=2}^n \frac{4a_i A_i W_0}{\mathcal{V}^2} \tau_i e^{-a_i \tau_i} + \frac{3\hat{\xi} W_0^2}{4\mathcal{V}^3} + \frac{D}{\mathcal{V}^\gamma}$$

$$m_{\tau_i}^2 \simeq \frac{W_0^2 (\ln \mathcal{V})^2 M_{\text{pl}}^2}{\mathcal{V}^2} \gg m_{\mathcal{V}}^2 \simeq \frac{W_0^2 M_{\text{pl}}^2}{\mathcal{V}^3 \ln \mathcal{V}}$$

taking all small moduli at minima: two field potential

$$V_{\text{inf}} = -\frac{3W_0^2}{2\mathcal{V}^3} \left(\sum_{i=2}^{n-1} \left[\frac{\lambda_i \alpha}{a_i^{3/2}} \right] (\ln \mathcal{V})^{3/2} - \frac{\xi}{2} \right) + \frac{D}{\mathcal{V}^\gamma} - \frac{4a_n A_n W_0}{\mathcal{V}^2} \tau_n e^{-a_n \tau_n}$$

modulus

inflaton

Shift in Volume Modulus

Potential experienced by the volume modulus depends on the inflaton: Vacuum misalignment

Other modulus are not shifted and having masses much larger than the Hubble scale.

$$V = -\frac{3W_0^2}{2\mathcal{V}^3} \left(\sum_{i=2}^n \left[\frac{\lambda_i \alpha}{a_i^{3/2}} \right] (\ln \mathcal{V})^{3/2} - \frac{\hat{\xi}}{2} \right) + \frac{D}{\mathcal{V}^\gamma}$$

$$Y = \frac{\delta\varphi}{M_{pl}} = \sqrt{\frac{2}{3}} \delta\phi \simeq 2\sqrt{\frac{2}{3}} R\phi_* \simeq 0.1 - 1$$

We validate the generic arguments with explicit calculations

e-foldings

t_1 : end of inflation

$$\rho_{\tau_n}(t_1) \approx \frac{M_{\text{pl}}^4 W_0^2 \beta}{\mathcal{V}^3}$$

Barnaby, Bond, Huang, Kofman (2009)

$$H(t_1) \approx \frac{M_{\text{pl}} W_0 \beta^{1/2}}{\mathcal{V}^{3/2}} \simeq m_{\mathcal{V}}$$

Volume modulus starts to oscillate immediately at the end of inflation

$$\rho_{\mathcal{V}}(t_1) \approx m_{\mathcal{V}}^2 \varphi_{\text{in}}^2 \approx \frac{M_{\text{pl}}^4 W_0^2 Y^2}{\mathcal{V}^3 \ln \mathcal{V}} \quad \frac{\rho_{\mathcal{V}}(t_1)}{\rho_{\tau_n}(t_1)} \approx \frac{Y^2}{\beta \ln \mathcal{V}} \equiv \theta^2 \ll 1$$

short matter dominated epoch until inflaton decays

$$N_{\text{mod1}} = \ln \left(\frac{a(t_2)}{a(t_1)} \right) = \frac{1}{3} \ln \left(\frac{\rho_{\tau_n}(t_1)}{\rho_{\tau_n}(t_2)} \right) \simeq \frac{2}{3} \ln \left(\frac{H(t_1)}{\Gamma_{\tau_n}} \right) \simeq \frac{2}{3} \ln \left(\frac{10 \beta^{1/2} \mathcal{V}^{1/2}}{W_0^2 (\ln \mathcal{V})^3} \right)$$

t_2 : inflation decay time

e-foldings

Inflaton may decay to hidden sectors Cicoli, Mazumdar (2010)

When inflaton decays, radiation domination

$$H(t_2) = H(t_1) \left(\frac{a(t_1)}{a(t_2)} \right)^{3/2} = H(t_1) e^{-\frac{3}{2} N_{\text{mod}1}} \simeq \frac{H(t_1) W_0^2 (\ln \mathcal{V})^3}{10 \beta^{1/2} \mathcal{V}^{1/2}}$$

At t_{eq} , radiation and volume modulus oscillations density become equal

$$\rho_{\text{rad}}(t_2) \left(\frac{a(t_2)}{a(t_{\text{eq}})} \right)^4 = \rho_{\mathcal{V}}(t_2) \left(\frac{a(t_2)}{a(t_{\text{eq}})} \right)^3 \quad H(t_{\text{eq}}) = \frac{H(t_1) W_0^2 (\ln \mathcal{V})^3 \theta^4}{10 \beta^{1/2} \mathcal{V}^{1/2}}$$

$$N_{\text{mod}2} \simeq \frac{2}{3} \ln \left(\frac{H(t_{\text{eq}})}{\Gamma_{\mathcal{V}}} \right)$$

Similar to the previous expressions

$$N_{\text{mod}2} \approx \frac{2}{3} \ln \left(\frac{16\pi \mathcal{V}^{5/2} (\ln \mathcal{V})^{5/2} Y^4}{10\beta^2} \right) \approx \frac{2}{3} \ln \left(\frac{16\pi \mathcal{V}^{5/2} Y^4}{10P^2 R^2 (\ln \mathcal{V})^{1/2}} \right)$$

A benchmark example

$$W_0 = \alpha = \lambda_i = 1, a_i = 2\pi, g_s = 0.06$$

$$\mathcal{V}_{\text{in}} \simeq 1.38 \cdot 10^5, \beta \simeq 3.88$$

$$\tau_n \simeq 1.12 \cdot 10^8 N_e^{-4} \quad \text{and} \quad r = 16\epsilon \simeq 1.04 \cdot 10^{-10}$$

plays important role



$$m_\nu \sim 10^8 - 10^9 \text{ GeV}$$

$$N_{\text{mod}1} \simeq 0.99 \quad \text{and} \quad N_{\text{mod}2} \simeq 25.4$$

$$N_e \simeq 44.65 + \frac{1}{4} \ln \left(\frac{\rho_*}{\rho_{\text{end}}} \right) \simeq 45 \quad \Rightarrow \quad \tau_n \simeq 27.3 \quad \text{and} \quad n_s \simeq 0.955$$

$$T_{\text{rh}} \gtrsim 10^3 \text{ GeV} \quad \text{OK with BBN bound}$$

Predictions for Kahler
Moduli Inflation shifts!



Yesterday's talk

Starobinsky-like Inflation, Supercosmology
and Neutrino Masses in No-Scale Flipped SU(5)

1704.07331

John Ellis^a, Marcos A. G. Garcia^b, Natsumi Nagata^c,
Dimitri V. Nanopoulos^d and Keith A. Olive^e

Entropy production from a non-inflaton field

$$N_* = N_*^{\text{STH}} - \frac{1}{3} \ln \Delta$$

For 'weak reheating', the spectral-index prediction is
changed substantially!

Post-inflationary history is crucial to understand!

Conclusions

- modulus dominated cosmology is a generic feature of string/sugra motivated scenario

Conclusions

- modulus dominated cosmology is a generic feature of string/sugra motivated scenario

$$\hat{N}_{inf} = 55 - \frac{1}{3} \left(\frac{\sqrt{16\pi} M_{Pl} Y^2}{m_\varphi} \right) \leftarrow$$

sensitivity

$$n_s \sim 10^{-3}$$

CORE/CMB IV

Conclusions


- modulus dominated cosmology is a generic feature of string/sugra motivated scenario

$$\hat{N}_{inf} = 55 - \frac{1}{3} \left(\frac{\sqrt{16\pi} M_{Pl} Y^2}{m_\varphi} \right) \simeq 45$$



- Independent constraint on modulus mass derived using precision CMB data

Bound..



for $N > 44$,
the bound
much
stronger than
BBN bound

Conclusions

- modulus dominated cosmology is a generic feature of string/sugra motivated scenario

$$\hat{N}_{inf} = 55 - \frac{1}{3} \left(\frac{\sqrt{16\pi} M_{Pl} Y^2}{m_\varphi} \right) \simeq 45$$



- Independent constraint on modulus mass derived using precision CMB data
- Explicit calculations for Kahler Moduli inflation

Thank You

Shift in Volume Modulus

Potential experienced by the volume modulus depends on the inflaton: Vacuum misalignment

Other modulus are not shifted and having masses much larger than the Hubble scale.

$$V = -\frac{3W_0^2}{2\mathcal{V}^3} \left(\sum_{i=2}^n \left[\frac{\lambda_i \alpha}{a_i^{3/2}} \right] (\ln \mathcal{V})^{3/2} - \frac{\hat{\xi}}{2} \right) + \frac{D}{\mathcal{V}^\gamma}$$

$$V(\mathcal{V}_*) = \frac{\partial V(\mathcal{V}_*)}{\partial \mathcal{V}} = 0 \quad -\frac{3W_0^2}{2} e^{-3\phi_*} \left(P \phi_*^{3/2} - \frac{\hat{\xi}}{2} \right) + D e^{-2\phi_*} = 0$$

$$\frac{3W_0^2}{2} e^{-3\phi_*} \left(3P \phi_*^{3/2} - \frac{3}{2} P \phi_*^{1/2} - \frac{3\hat{\xi}}{2} \right) - 2D e^{-2\phi_*} = 0$$

$$\phi \equiv \ln \mathcal{V} \quad \text{and} \quad P \equiv \alpha \sum_{i=2}^n \lambda_i a_i^{-3/2} = \frac{\alpha}{R} \lambda_n a_n^{-3/2} \quad R \equiv \frac{\lambda_n a_n^{-3/2}}{\sum_{i=2}^n \lambda_i a_i^{-3/2}} \ll 1$$

Shift in Volume Modulus

$$\phi_*^{3/2} - \frac{3}{2}\phi_*^{1/2} - \frac{\hat{\xi}}{2P} = 0$$

$$D = \frac{9W_0^2}{4} P e^{-\phi_*} \phi_*^{1/2}$$

$$V_{\text{in}}(\phi) = -\frac{3W_0^2}{4} e^{-3\phi} \left[2P(1-R)\phi^{3/2} - \hat{\xi} - 3P\phi_*^{1/2} e^{(\phi-\phi_*)} \right]$$

$$(1-R)\phi_{\text{in}}^{3/2} - \frac{1}{2}(1-R)\phi_{\text{in}}^{1/2} - e^{(\phi_{\text{in}}-\phi_*)}\phi_*^{1/2} - \frac{\hat{\xi}}{2P} = 0$$

$$V_{\text{in}}(\phi) = V(\phi) + \delta V(\phi)$$

$$\delta V(\phi) = \frac{3W_0^2}{2} e^{-3\phi} P R \phi^{3/2}$$

$$\delta\phi = -\frac{\delta V'(\phi_*)}{V''(\phi_*)} = 4R \frac{\phi_* + \frac{\hat{\xi}}{2P} \phi_*^{1/2}}{2\phi_* - 1} \simeq 2R\phi_*$$

$$\delta\phi = \phi_{\text{in}} - \phi_*$$

$$Y = \frac{\delta\varphi}{M_{pl}} = \sqrt{\frac{2}{3}}\delta\phi \simeq 2\sqrt{\frac{2}{3}}R\phi_* \simeq 0.1 - 1$$

We validate the generic arguments with explicit calculations

Inflationary Phenomenology

$$V = V_0 - \frac{4W_0 a_n A_n}{\mathcal{V}_{\text{in}}^2} \left(\frac{3\mathcal{V}_{\text{in}}}{4\lambda} \right)^{2/3} \sigma^{4/3} \exp \left[-a_n \left(\frac{3\mathcal{V}_{\text{in}}}{4\lambda} \right)^{2/3} \sigma^{4/3} \right]$$

Canonical inflaton field $V(\sigma) = C_0(1 - e^{-b\sigma})$

Effective single field dynamics

Volume modulus is stabilised during inflation and heavy

$$\epsilon = \frac{M_{\text{pl}}^2}{2} \left(\frac{V'}{V} \right)^2 = \frac{32\mathcal{V}_{\text{in}}^3}{3\beta^2 W_0^2 \lambda_n} a_n^2 A_n^2 \sqrt{\tau_n} (1 - a_n \tau_n)^2 e^{-2a_n \tau_n},$$

$$\eta = M_{\text{pl}}^2 \frac{V''}{V} = -\frac{4\mathcal{V}_{\text{in}}^2}{3\beta W_0 \lambda_n \sqrt{\tau_n}} a_n A_n \left[(1 - 9a_n \tau_n + 4a_n^2 \tau_n^2) e^{-a_n \tau_n} \right].$$

Inflationary Phenomenology

$$N_e(\sigma) = \int_{\sigma_{\text{end}}}^{\sigma} \frac{1}{\sqrt{2\epsilon(\sigma)}} d\sigma \simeq \frac{3\beta W_0 \lambda_n}{16 \mathcal{V}_{\text{in}}^2 a_n^{3/2} A_n} \frac{e^{a_n \tau_n}}{(a_n \tau_n)^{3/2}}$$

$$\epsilon \simeq \left(\frac{3\lambda_n}{8a_n^{3/2} \mathcal{V}_{\text{in}}} \right) \frac{1}{N_e^2 \sqrt{a_n \tau_n}} \ll \eta \simeq -\frac{1}{N_e}$$

COBE normalisation: $\frac{V^{3/2}}{M_{\text{pl}}^3 V'} = 5.2 \times 10^{-4}$

$$\tau_n \simeq 7.31 \cdot 10^{-14} \left(\frac{6\pi \lambda_n}{g_s \beta e^{K_{\text{cs}}}} \right)^2 \left(\frac{\mathcal{V}_{\text{in}}^4}{W_0^4 a_n^4} \right) \frac{1}{N_e^4}$$

$$r = 16\epsilon \simeq 16 \times 3.7 \cdot 10^6 \left(\frac{g_s \beta e^{K_{\text{cs}}}}{16\pi} \right) \left(\frac{W_0^2}{\mathcal{V}_{\text{in}}^3} \right) \ll 10^{-4} N_e^{-3}$$

$$n_s = 1 + 2\eta - 6\epsilon \simeq 1 - \frac{2}{N_e}$$

spectral index depends only on N_e