

Matter Wave Ramsey Interferometry & The Quantum Nature of Gravity


Sougato Bose

University College London

Based on:

- M. Scala, M. S. Kim, G. W. Morley, P. F. Barker, S. Bose, Phys. Rev. Lett. **111**, 180403 (2013).
- C. Wan, M. Scala, G. W. Morley, A. Rahman, H. Ulbricht, J. Bateman, P. F. Baker, S. Bose, M. S. Kim, Phys. Rev. Lett. **117**, 143003 (2016).
- S. Bose, A. Mazumdar, G. W. Morley, H. Ulbricht, M. Paternostro, P. F. Barker, A. Geraci, M. S. Kim, G. J. Milburn, arXiv (soon)


***Tiny* Superpositions of a Macroscopic Object** (the older idea was to investigate through the decoherence induced into the ancilla by the macroscopic object)



S. Bose, K. Jacobs, P. L. Knight,
Phys. Rev. A 59 (5), 3204
(1999).

+

Armour, Blencowe, Schwab,
PRL 2002.




Marshall, Simon, Penrose,
Bouwmeester, PRL 2003.

Bose, PRL 2006.

Ramsey Interferometry with a Levitated Thermal Mesoscopic Object

Diamond bead trapped in an optical trap. The bead contains a spin-1 NV center.



Coupling between the spin and the motion

Calling (0,0,0) the position of the minimum of the potential, let us consider the magnetic field of a magnetized sphere with magnetic dipole $\mathbf{m}=(0,0,m_z)$ placed at the position (0, 0, z_0). Expanding it up to first order around the center of the trap, the magnetic field is:

$$B_x = -\frac{3\mu_0 m_z}{4\pi|z_0|^4} \cdot \frac{z_0}{|z_0|} x,$$

$$B_y = -\frac{3\mu_0 m_z}{4\pi|z_0|^4} \cdot \frac{z_0}{|z_0|} y,$$

$$B_z = \frac{\mu_0 m_z}{2\pi |z_0|^3} + \frac{3\mu_0 m_z}{2\pi|z_0|^4} \cdot \frac{z_0}{|z_0|} z.$$

The zeroeth order term in B gives a Zeeman splitting between $|+1\rangle$ and $|-1\rangle$.

Coupling between the spin and the motion

The linear term in the expansion gives the following coupling between the spin and the vibrational motion:

$$H_{\text{int}} = -\lambda[2 S_z (c + c^\dagger) - A_x S_x (a + a^\dagger) - A_y S_y (b + b^\dagger)],$$

where
:

$$A_x = \sqrt{\frac{\omega_z}{\omega_x}}, \quad A_y = \sqrt{\frac{\omega_z}{\omega_y}},$$

and:

$$\lambda = \frac{3\mu_0 m_z z_0}{4\pi |z_0|^5} g_{NV} \mu_B \sqrt{\frac{\hbar}{2m\omega_z}}$$

Spin Optomechanical coupling also derived in:


Z Yin, T Li, X Zhang, LM Duan, Physical Review A 88, 033614 (2013).

Also Rabl et. al. (2008), Oosterkamp et. al. (2008)

Spin-dependent displacement in a gravitational field

In the limit case of infinite ω_x and ω_y the Hamiltonian describes a conditional displacement of the trapping potential, whose direction depends on the value of S_z :

$$H = DS_z^2 + \hbar\omega_z c^\dagger c - 2\lambda S_z(c + c^\dagger) + 2\Delta\lambda(c + c^\dagger)$$



The interferometric scheme

Suppose that the oscillator is initially in a coherent state $|\beta\rangle$ and that the spin is in the eigenstate $|S_z=0\rangle$:


$$|\Psi(0)\rangle = |\beta\rangle |0\rangle$$

Step 1: apply a very rapid mw pulse which transforms the state of the spin according to:

$$|0\rangle \rightarrow \frac{1}{\sqrt{2}} (|+1\rangle + |-1\rangle)$$

So that we obtain the superposition of two states which oscillate in opposite directions.

Evolution of an *arbitrary* coherent state
(at the time period T it comes back!)



$$|\beta\rangle|\pm 1\rangle \rightarrow e^{-\frac{i}{\hbar}\left(D - \frac{3(\lambda - \Delta\lambda)^2}{\hbar\omega_Z}\right)t} e^{-\frac{i}{\hbar}\frac{(\lambda \mp \Delta\lambda)^2}{\hbar\omega_Z^2}\sin\omega_Z t} \left| \left(\beta \mp \frac{\lambda \mp \Delta\lambda}{\hbar\omega_Z} \right) e^{-i\omega_Z t} \pm \frac{\lambda \mp \Delta\lambda}{\hbar\omega_Z} \right\rangle |\pm 1\rangle$$

Evolution

Step 2: the state evolves for time T equal to the period of the oscillator; the oscillation is different according to the spin, but at $t=T$ the vibrational state is the same for both $|+1\rangle$ and $|-1\rangle$

so that the spin state is separated from the vibrational state and is (up to a global phase):

$$|\Psi_{spin}\rangle = \frac{1}{\sqrt{2}} \left(|+1\rangle + e^{i\Delta\phi} |-1\rangle \right)$$

with:

$$\Delta\phi = -\frac{12\lambda \Delta\lambda}{\hbar^2 \omega_z} T$$

$m = 10^{-17}$ Kg, $\omega_z = 100$ kHz, $\Delta x = 1$ pm, $\Delta t \sim 1$ μ s, Gradient $\sim 10^4$ T/m, we have **$\Delta\phi \sim 1$**

Measuring the phase shift due to gravitational potential difference

Step 3: apply the same very rapid mw pulse as in step 1, which, in the absence of the phase difference $\Delta\phi$ transforms the state of the spin according to:

$$\frac{1}{\sqrt{2}} (|+1\rangle + |-1\rangle) \rightarrow |0\rangle$$

The presence of $\Delta\phi$ gives a modulation of the population of $|S_z=0\rangle$ according to:


$$P_0(T + \Delta t_{pulse}) = \cos^2 \frac{\Delta\phi}{2}$$

Free particle in an inhomogeneous magnetic field (acceleration $+a$ or $-a$)


$$x_{\sigma}(t,j)=x_j(0)\pm\frac{1}{2}at^2$$

$$=\frac{a\tau}{4}(t-\frac{\tau}{4})\mp\frac{1}{2}a(t-\frac{\tau}{4})^2$$

$$=\frac{1}{2}a(\frac{\tau}{4})^2\mp\frac{a\tau}{4}(t-\frac{3\tau}{4})\pm\frac{1}{2}a(t-\frac{3\tau}{4})^2$$



Free flight scheme able to achieve 100 nm separation among superposed components:



$$|\Psi(t_3)\rangle = \frac{1}{\sqrt{2}}|\psi(t_3)\rangle(|+1\rangle + e^{-i\phi_g}|-1\rangle)$$


$$\langle x|\psi(t_3)\rangle = e^{-ip_0x}e^{-[(x-x_0-p_0t_3/m-g\cos\theta t_3^2/2)^2/2(\sigma')^2]}$$

$$\phi_g = (1/16\hbar)gt_3^3g_{\rm NV}\mu_B(\partial B/\partial x)\cos\theta$$


$$\Delta x_M = 2 \times \frac{1}{2m} g_{\rm NV} \mu_B \frac{\partial B}{\partial x} (t_3/4)^2$$

10^{10} amu mass can be placed in a superposition of states separated by 100 nm.

A Schematic of two matter-wave interferometers near each other




Consider two neutral test masses *held* in a superposition, each exactly as a path encoded qubit (states $|L\rangle$ and $|R\rangle$), near each other.



where


$$\phi_{RL} \sim \frac{Gm_1m_2\tau}{\hbar(d - \Delta x)}, \phi_{LR} \sim \frac{Gm_1m_2\tau}{\hbar(d + \Delta x)},$$

$$\phi_{LL} = \phi_{RR} \sim \frac{Gm_1m_2\tau}{\hbar d}$$



$$\begin{aligned}
 |\Psi(t = \tau)\rangle_{12} &= \frac{1}{2} (e^{i\phi_{LL}} |L\rangle_1 |L\rangle_2 + e^{i\phi_{LR}} |L\rangle_1 |R\rangle_2 \\
 &\quad + e^{i\phi_{RL}} |R\rangle_1 |L\rangle_2 + e^{i\phi_{RR}} |R\rangle_1 |R\rangle_2) \\
 &= \frac{e^{i\phi_{RR}}}{\sqrt{2}} \left\{ |L\rangle_1 \frac{1}{\sqrt{2}} (|L\rangle_2 + e^{i\Delta\phi_{LR}} |R\rangle_2) \right. \\
 &\quad \left. + |R\rangle_1 \frac{1}{\sqrt{2}} (e^{i\Delta\phi_{RL}} |L\rangle_2 + |R\rangle_2) \right\}
 \end{aligned}$$

The above state is maximally entangled when $\Delta\phi_{LR} + \Delta\phi_{RL} \sim \pi$.



For

$$d - \Delta x \ll d, \Delta x,$$

we have

$$\Delta\phi_{RL} \sim \frac{Gm_1m_2\tau}{\hbar(d - \Delta x)} \gg \Delta\phi_{LR}, \Delta\phi_{LL}, \Delta\phi_{RR}$$


For


$$d - \Delta x \ll d, \Delta x,$$

we have

$$\Delta\phi_{RL} \sim \frac{Gm_1m_2\tau}{\hbar(d - \Delta x)} \gg \Delta\phi_{LR}, \Delta\phi_{LL}, \Delta\phi_{RR}$$

For mass $\sim 10^{-14}$ kg (microspheres), separation at closest approach of the masses ~ 100 microns (to prevent Casimir interaction), time ~ 1 seconds, $\Delta\phi_{RL} \sim 1$





Spin Correlation Functions Certifying Entanglement

What does it imply in the context of low energy effective field theory?

$$\mathcal{H} = \sum_{j,\sigma} m_j c^2 a_{\sigma,j}^\dagger a_{\sigma,j} + \sum_{\mathbf{k}} \hbar \omega_k b_{\mathbf{k}}^\dagger b_{\mathbf{k}} - \hbar \sum_{j,\mathbf{k},\sigma} g_{j,\mathbf{k}} a_{\sigma,j}^\dagger a_{\sigma,j} (b_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}_{j,\sigma,t}} + b_{\mathbf{k}}^\dagger e^{-i\mathbf{k} \cdot \mathbf{r}_{j,\sigma,t}})$$

Blencowe,
PRL 2013

$$g_{j,\mathbf{k}} = m_j c^2 \sqrt{\frac{8\pi G}{\hbar c^3 k V}}$$

$$|\Psi_{\text{mat+grav}}(t)\rangle = \frac{1}{2} \sum_{\sigma,\sigma'} a_{1,\sigma}^\dagger a_{2,\sigma'}^\dagger |0\rangle$$

$$\otimes \prod_{\mathbf{k}} e^{i \frac{|g_{1,\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}_{1,\sigma,t}} + g_{2,\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}_{2,\sigma',t}}|^2}{\omega_k} t} |\alpha_{\mathbf{k},\sigma,\sigma'}\rangle_{\mathbf{k}}$$

$$\alpha_{\mathbf{k},\sigma,\sigma'} = \left(\frac{g_{1,\mathbf{k}}}{\omega_k} e^{i\mathbf{k} \cdot \mathbf{r}_{1,\sigma,t}} + \frac{g_{2,\mathbf{k}}}{\omega_k} e^{i\mathbf{k} \cdot \mathbf{r}_{2,\sigma',t}} \right) (e^{i\omega_k t} - 1)$$

Superpositions of *distinct* (?) coherent states of the gravitational field

Conclusions

Long term motive: Enhancing the domain of investigation of the quantum superposition principle

M. Scala, M. S. Kim, G. W. Morley, P. F. Barker, S. Bose, Phys. Rev. Lett. **111**, 180403 (2013)
Free flight scheme: C. Wan et. al., Phys. Rev. Lett. **117**, 143003 (2016).

Long term motive: To test the quantum nature of gravity.
S. Bose, A. Mazumdar, G. W. Morley, H. Ulbricht, M. Paternostro,
P. F. Barker, A. Geraci, M. S. Kim, G. J. Milburn, arXiv (soon)