


Diffusion versus Superdiffusion in a stochastic Hamiltonian lattice field model

Cédric Bernardin

(collaborators: P. Gonçalves, M. Jara, M. Simon)

University of Nice, France



- *One dimensional conservative asymmetric* interacting particle systems display anomalous diffusion.
- It means that if the system at equilibrium is locally disturbed by adding some extra energy, the perturbation will not diffuse like a Brownian Motion but like some *superdiffusive process*.
- What is the superdiffusive process? How *universal* is it?

- We are interested in one dimensional interacting particle systems which conserve some quantities (energy, density, momentum ...).
- In a suitable *space-time scale*, the empirical conserved quantities (macroscopic, coarse-grained) will evolve according to some hyperbolic system of conservation laws (e.g. Euler equations). These are the *hydrodynamic limits* of the system.
- Starting from these macroscopic equations (ignoring the details of the microscopic dynamics), the *Nonlinear Fluctuating Hydrodynamics Theory* (Spohn) predicts very precisely the form of the fluctuations of the conserved quantities.

Microscopic models

- We consider a lattice field model $\{\eta_x(t) \in \mathbb{R}; x \in \mathbb{Z}\}$ whose dynamics is composed of a deterministic part and of a stochastic part [B. Stoltz'11], [Basile, B., Olla'06] .
- The stochastic part is introduced to provide a better control of the chaotic motion due to the nonlinearities of the interactions.

- **Deterministic part:** It is given by Hamilton equations:

$$d\eta_x = (V'(\eta_{x+1}) - V'(\eta_{x-1})) dt, \quad x \in \mathbb{Z},$$

where $V : \mathbb{R} \rightarrow \mathbb{R}$ is a well behaved potential.


- **Deterministic part:** It is given by Hamilton equations:

$$d\eta_x = (V'(\eta_{x+1}) - V'(\eta_{x-1})) dt, \quad x \in \mathbb{Z},$$

where $V : \mathbb{R} \rightarrow \mathbb{R}$ is a well behaved potential.

- **Stochastic part :**

Independent Poisson processes (clock) on each bond $\{x, x + 1\}$. When the clock of $\{x, x + 1\}$ rings, η_x is exchanged with η_{x+1} . The dynamics between two successive rings of the clocks is given by the Hamiltonian dynamics.



Conserved quantities (slow variables):

Conserved quantities (slow variables):

1. The energy $\sum_x e_x = \sum_x V(\eta_x)$.

Conserved quantities (slow variables):

1. The energy $\sum_x e_x = \sum_x V(\eta_x)$.
2. The volume $\sum_x \eta_x$.

- The energy $\sum_x V(\eta_x)$ and the volume $\sum_x \eta_x$ are the **only** conserved quantities of the stochastic model (in a suitable sense) [B. Stoltz'11], [Fritz-Funaki-Lebowitz'93].
- The Gibbs equilibrium measures $\langle \cdot \rangle_{\tau, \beta}$ are parameterized by two parameters $(\tau, \beta) \in \mathbb{R} \times [0, \infty)$ and are product

$$\langle \cdot \rangle_{\tau, \beta} \sim \exp\{-\beta \sum_x (V(\eta_x) + \tau \eta_x)\} d\eta$$

are invariant for the dynamics.

Hydrodynamics: Euler equations

Theorem (B., Stoltz'11)

For $t < T^$ (first shock), the hydrodynamic equations of the stochastic dynamics are given by the compressible Euler equations:*

$$\begin{cases} \partial_t \mathbf{v} = 2\partial_q \mathcal{P}, \\ \partial_t \mathbf{e} = \partial_q \mathcal{P}^2. \end{cases}$$

In the harmonic case $V(r) = r^2/2$, $\mathcal{P}(\mathbf{v}, \mathbf{e}) = \mathbf{v}$ and $T^ = \infty$ (no shocks).*

Proof based on [Olla, Varadhan, Yau'91], [Fritz-Funaki-Lebowitz'93].

Nonlinear fluctuating hydrodynamics predictions

The theory of *nonlinear fluctuating hydrodynamics* (Spohn) predicts the long time behavior of the equilibrium time-space correlation functions of the conserved fields $g(x, t) = (\eta_x(t), e_x(t))$

$$S_{\alpha\alpha'}(x, t) = \langle g_\alpha(x, t) g_{\alpha'}(0, 0) \rangle_{\tau, \beta} - \langle g_\alpha \rangle_{\tau, \beta} \langle g_{\alpha'} \rangle_{\tau, \beta}$$

Gibbs measure: $\langle \cdot \rangle_{\tau, \beta} \sim \exp\left\{-\beta \sum_x (e_x + \tau \eta_x)\right\} d\eta.$

temperature: β^{-1} pressure: τ

- Spohn's theory: the long time behavior of the correlation functions of the conserved fields depends only on the function $(v, e) \rightarrow \mathcal{P}(v, e)$, and parameters τ, β , but **NOT** on the details of the microscopic dynamics.
- It is a *macroscopic* theory based on the validity of the hydrodynamics in the Euler time scale.

- $n = 1$ case: 2 UC
 - Edwards-Wilkinson (Gaussian)
 - "KPZ fixed point" (non Gaussian) ^a
- $n \geq 2$ case:
 - richer (many UC),
 - different time scales involved

^ascaling limit of the solution of the KPZ equation

$G^1 \backslash G^2$	$\begin{pmatrix} * \\ * \end{pmatrix}$	$\begin{pmatrix} * \\ 0 \end{pmatrix}$	$\begin{pmatrix} 0 \\ * \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$
$\begin{pmatrix} * \\ * \end{pmatrix}$	3/2	3/2	5/3	2
$\begin{pmatrix} 0 \\ * \end{pmatrix}$	3/2	3/2	5/3	2
$\begin{pmatrix} * \\ 0 \end{pmatrix}$	3/2	3/2	$\frac{1+\sqrt{5}}{2}$	2
$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	2	2	2	2

Popkov et al., J. Stat. Phys. 160 (2015), $n = 2$

We consider now the harmonic case

$$V(r) = r^2/2 \Rightarrow e_x = \eta_x^2/2$$

Our results are in *partial agreement* with Spohn's predictions



Need some adjustments in the NFH

We define the space-time correlation of the energy

$$E_t(x) = \left\langle \left(e_0(0) - \frac{1}{\beta} \right) \left(e_t(x) - \frac{1}{\beta} \right) \right\rangle_{\tau, \beta}$$

and the space-time correlation of the volume

$$V_t(x) = \left\langle \left(\eta_0(0) - \tau \right) \left(\eta_t(x) - \tau \right) \right\rangle_{\tau, \beta}$$

Theorem (B., Gonçalves, Jara'16)

- Volume: $\lim_{n \rightarrow \infty} V_{tn^2}([nq]) = \frac{2}{\beta} \mathcal{V}_t(q), \quad t > 0, \quad q \in \mathbb{R},$

$$\partial_t \mathcal{V} = \Delta \mathcal{V}, \quad \text{heat equation}$$

- Energy: $\lim_{n \rightarrow \infty} E_{tn^{3/2}}([nq]) = \frac{2}{\beta^2} \mathcal{E}_t(q), \quad t > 0, \quad q \in \mathbb{R},$

$$\partial_t \mathcal{E} = -\frac{1}{\sqrt{2}} \{ (-\Delta)^{3/4} - \nabla (-\Delta)^{1/4} \} \mathcal{E}, \quad \text{skew fractional heat equation}$$

See also [Basile-B.-Olla'06,'09, Basile-Olla-Spohn'08, Mellet-Mischler-Mouhot'08, Jara-Komorowski-Olla'09, Delfini-Lepri-Livi-Mejia-Monasterio-Politi'10, Jara-Komorowski-Olla'15]

Proof:

We will relate the fractional Laplacian with solutions of the following extension problem. For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, we consider the extension $u : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}$ that satisfies the equation

$$u(x, 0) = f(x) \quad (1.4)$$

$$\Delta_x u + \frac{a}{y} u_y + u_{yy} = 0 \quad (1.5)$$

The equation (1.5) can also be written as

$$\operatorname{div}(y^a \nabla u) = 0 \quad (1.6)$$

Which is clearly the Euler-Lagrange equation for the functional

$$J(u) = \int_{y>0} |\nabla u|^2 y^a dX \quad (1.7)$$

We will show that

$$C(-\Delta)^s f = \lim_{y \rightarrow 0^+} -y^a u_y = \frac{1}{1-a} \lim_{y \rightarrow 0} \frac{u(x, y) - u(x, 0)}{y^{1-a}} \quad \text{with } s=(1-a)/2$$

Caffarelli & Silvestre, An extension problem related to the fractional Laplacian, CPDE '07

Weak anharmonicity limit: Universality persistence

Weak anharmonic potential : $V(r) = r^2 + \gamma_n r^4/4$.

Weak anharmonicity limit: Universality persistence

Weak anharmonic potential : $V(r) = r^2 + \gamma_n r^4/4$.

- If $\gamma_n = 0$ then the energy fluctuation field is described in the time scale $tn^{3/2}$ by a $3/2$ stable asymmetric Lévy process and the volume fluctuation field is described in the time scale tn^2 by a Brownian motion.

Weak anharmonicity limit: Universality persistence

Weak anharmonic potential : $V(r) = r^2 + \gamma_n r^4/4$.

- If $\gamma_n = 0$ then the energy fluctuation field is described in the time scale $tn^{3/2}$ by a $3/2$ stable asymmetric Lévy process and the volume fluctuation field is described in the time scale tn^2 by a Brownian motion.
- If $\gamma_n = O(1)$ and $\tau = 0, \beta > 0$, then NFH [Spohn-Stoltz'15] indicates that the energy-volume fluctuation fields are described by the same processes.

Theorem (B., Gonçalves, Jara, Simon '17)

The harmonic scenario persists in the nonlinear regime if γ_n is sufficiently small. It is valid for the energy for $\gamma_n \ll n^{-1/4}$ and for the volume for $\gamma_n \ll n^{-1/2}$.

Extension of this result outside of this regime requires new ideas.

Crossover between Diffusion and Superdiffusion

How can we cross different Universality Classes by tuning the parameters of the model ?

$G^2 \backslash G^1$	$\begin{pmatrix} * \\ * \end{pmatrix}$	$\begin{pmatrix} * \\ 0 \end{pmatrix}$	$\begin{pmatrix} 0 \\ * \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$
$\begin{pmatrix} * \\ * \end{pmatrix}$	$3/2$	$3/2$	$5/3$	2
$\begin{pmatrix} 0 \\ * \end{pmatrix}$	$3/2$	$3/2$	$5/3$	2
$\begin{pmatrix} * \\ 0 \end{pmatrix}$	$5/3$	$5/3$	$\frac{1+\sqrt{5}}{2}$	$3/2$
$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	2	2	2	2

Popkov et al., J. Stat. Phys. 160 (2015)

Weakly harmonic chain


- Consider the harmonic chain where the potential $V(r)$ is now

$$V(r) = \frac{c}{n^b} r^2$$

with $c, b > 0$ two positive constants.

- We look at the system in the time scale tn^a , $a > 0$, such that the energy field has a non-trivial limit.

[B., Gonçalves, Jara'16]



For $b = 1/3$, $a = 2$ the **energy limiting field** is described by a Levy process interpolating between the asymmetric stable Levy process and the Brownian motion.

- The generator of the interpolating Levy process is

$$\Delta - \frac{c^{3/2}}{\sqrt{2}} \{(-\Delta)^{3/4} - \nabla(-\Delta)^{1/4}\}.$$

- As $c \rightarrow \infty$, scaled with c , it goes to the skew 3/4-fractional Laplacian.
- As $c \rightarrow 0$, it goes to the Laplacian.

Evanescent flip noise limit

Evanescent flip noise limit

- Consider the initial process (harmonic chain + exchange noise) and add a second stochastic perturbation with intensity $\gamma_n = cn^{-b}$, $c, b > 0$, which consists to flip independently on each site at exponential times the variable η_x into $-\eta_x$.


Evanescent flip noise limit

- Consider the initial process (harmonic chain + exchange noise) and add a second stochastic perturbation with intensity $\gamma_n = cn^{-b}$, $c, b > 0$, which consists to flip independently on each site at exponential times the variable η_x into $-\eta_x$.
- The energy is conserved but the volume $\sum_x \eta_x$ is not (stricto sensu, only if $b = \infty$).

Evanescent flip noise limit

- Consider the initial process (harmonic chain + exchange noise) and add a second stochastic perturbation with intensity $\gamma_n = cn^{-b}$, $c, b > 0$, which consists to flip independently on each site at exponential times the variable η_x into $-\eta_x$.
- The energy is conserved but the volume $\sum_x \eta_x$ is not (stricto sensu, only if $b = \infty$).
- We look at the system in the time scale tn^a , $a > 0$, such that the energy field has a non-trivial limit.

[B., Gonçalves, Jara, Sasada, Simon '15], [B., Gonçalves, Jara, Simon '16]



For $b = 1$, $a = 3/2$ the **energy limiting field** is described by a Levy process interpolating between the Brownian motion and the asymmetric stable Levy process.

- The Fourier symbol of the generator of the interpolating Levy process is

$$\frac{1}{2\sqrt{3}} \frac{(2i\pi k)^2}{\sqrt{c + i\pi k}}, \quad k \in \mathbb{R}.$$

- As $c \rightarrow 0$ it goes to the Fourier symbol of the skew 3/4-fractional Laplacian.
- As $c \rightarrow \infty$, scaled with c , it goes to the Fourier symbol of the Laplacian.

Space of IIC

