## PROBLEM SET 4

July 18, 2019

- (1) Let X be a compact complex manifold with a Kähler metric  $g, E \to X$  a holomorphic vector bundle, and h Hermitian metric on E. Assume  $\Theta_g(h) \ge cI$  for some c > 0.
  - (a) Show that for any  $\phi \in L^2_{n,q}(g,h)$ , there exists  $v \in \Gamma(X, \wedge_X^{n,q} \otimes E)$  such that  $\overline{\partial}v$ ,  $\overline{\partial}^*v \in L^2_{n,q}(g,h)$  and  $\Box v = \phi$  in the weak sense, i.e., for all  $\eta \in \Gamma(X, \wedge_X^{n,q} \otimes E)$  smooth,

 $(\overline{\partial}v, \overline{\partial}\eta) + (\overline{\partial}^*v, \overline{\partial}^*\eta) = (\phi, \eta).$ 

- (b) Show that if  $\overline{\partial}\phi = 0$  then  $\overline{\partial}^*\overline{\partial}v = 0$  in the weak sense,  $(\overline{\partial}v, \overline{\partial}\eta) = 0$  for smooth  $\eta$ .
- (c) From (b), prove Hörmander for compact complex manifolds.
- (2) Show that if X is a compact complex manifold and  $L \to X$  is a holomorphic line bundle admitting a metric of positive curvature then
  - (a) X is Kähler, and
  - (b) Any  $\square$ -harmonic form is identically equal to 0.
- (3) Explani, using the complex geometry we have developed, the close relationship between dual connnections and formal adjoints.
- (4) Let X be a compact Kähler manifold,  $E \to X$  a holomorphic vector bundle with Hermitian metric h. Assume  $\Theta_g(h) \geq cI$ . Let  $\phi \in \Gamma(X, \wedge_X^{n,q} \otimes E)$  be a smooth form satisfying  $\overline{\partial} \phi = 0$ . Consider

$$\mathcal{A}_{\phi} := \{ \phi + \overline{\partial} \eta : \eta \in \Gamma(X, \wedge_X^{n, q} \otimes E) \}.$$

Find the element of  $\mathcal{A}_{\phi}$  whose  $L^2$ -norm is minimal.

(5) Let X be a compact complex manifold and  $L \to X$  a holomorphic line bundle with a Hermitian metric h whose curvature is positive. Let  $x_1, \ldots, x_n \in X$  be distinct points and  $v_i \in L_{x_i}$ ,  $1 \le i \le n$  be such that  $h(v_i, v_i) = 1$ . Show that there exists m sufficiently large and a holomorphic section s of  $L^{\otimes m} \to X$  such that  $s(x_i) = v_i$ ,  $1 \le i \le n$ .

1