
Out of equilibrium dynamics of complex systems

Leticia F. Cugliandolo

Sorbonne Université

Laboratoire de Physique Théorique et Hautes Energies

Institut Universitaire de France

`leticia@lpthe.jussieu.fr`

`www.lpthe.jussieu.fr/~leticia/seminars`

Bangalore, India, 2021

Plan of Lectures

1. Introduction
2. Coarsening
3. Active Matter
4. Disorder
5. Integrability

Second lecture

Plan of Lectures

1. Introduction
2. **Coarsening**
3. Active Matter
4. Disorder
5. Integrability

References

— Phase ordering kinetics, critical dynamics & general

A. J. Bray, *Theory of phase ordering kinetics*, Adv. Phys. **43**, 357 (1994).

S. Puri, *Kinetics of Phase Transitions*, (Vinod Wadhawan, 2009).

L. F. Cugliandolo, *Topics in coarsening phenomena*, arXiv:0911.0771, Physica A 389, 4360 (2010).

F. Corberi & P. Politi eds., *Coarsening dynamics*, Comptes Rendus Physique **16** (2015).

P. C. Hohenberg and B. I. Halperin, *Theory of dynamic critical phenomena*, Rev. Mod. Phys. **49**, 435 (1977).

U. C. Tauber, *Critical Dynamics: A Field Theory Approach to Equilibrium and Non-Equilibrium Scaling Behavior* (Cambridge University Press, 2014).

P. Krapivsky, S. Redner and E. Ben-Naim, *A kinetic view of statistical physics* (Cambridge University Press, 2010).

M. Henkel and M. Pleimling, *Non-Equilibrium Phase Transitions Volume 2: Ageing and Dynamical Scaling Far from Equilibrium*, (Springer, 2010).

Plan of this lecture

1. The phenomenon
2. Theoretical setting
3. Critical and sub-critical quenches
4. Dynamic scaling
5. Dynamic universality classes
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. Mean-field models
9. Modern studies

Plan of this lecture

1. **The phenomenon**
2. Theoretical setting
3. Critical and sub-critical quenches
4. Dynamic scaling
5. Dynamic universality classes
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. Mean-field models
9. Modern studies

Phenomenon

The talk focuses on a very well-known example

Dynamics following a change of a control parameter

- If there is an equilibrium phase transition, the **equilibrium phases** are known on both sides of the transition.
i.e. the asymptotic state is known.
- For a purely dynamic problem, the **absorbing states** are known.
- The **dynamic mechanism** towards equilibrium (or the absorbing states) is understood : the systems try to order locally in one of the few competing states.

Interests and goals

Practical interest, *e.g.*

- Mesoscopic structure effects on the opto-mechanical properties of phase separating glasses
- Cooling rate effects on the density of topological defects in cosmology and condensed matter

Fundamental interest, *e.g.*

- A theoretical problem beyond perturbation theory.
- Are there growth phenomena in problems with yet unknown dynamic mechanisms ? **e.g. glasses**
- Generic features of macroscopic systems out of equilibrium ?


Context

Open systems

Our interest is to describe the **dynamics** of a **classical** (or quantum) **system** coupled to a **classical** (or quantum) **environment**.

The Hamiltonian of the ensemble is

$$H = H_{syst} + H_{env} + H_{int}$$




The dynamics of all variables are given by **Newton** (or Heisenberg) rules, depending on the variables being classical (or quantum).

$$\mathcal{E}_{syst}(t) \neq \text{ct}, \text{ and } e_0 \ll \mathcal{E}_{syst} \ll \mathcal{E}_{env}.$$

$2d$ Ising model

Snapshots after an instantaneous quench from $T \rightarrow \infty$ to $T < T_c$




At $T = T_c$ **critical dynamics**

At $T < T_c$ **coarsening**

A certain number of **interfaces** or **domain walls** in the last snapshots.

Membranes Proteins


Phase separation



Wadsten, Wöhri, Snijder, Katona, Gardiner, Cogdell, Neutze, Engström,
Lipidic Sponge Phase Crystallization of Membrane Proteins, J. Mol. Biol. 06

Glasses


Phase separation



$t = 1 \text{ min}$

Glasses


Phase separation



$t = 4 \text{ min}$

Glasses


Phase separation



$t = 16 \text{ min}$

Glasses

Phase separation



$t = 64 \text{ min}$

Plan of this lecture

1. The phenomenon
2. **Theoretical setting**
3. Critical and sub-critical quenches
4. Dynamic scaling
5. Dynamic universality classes
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. Mean-field models
9. Modern studies

Kinetic Ising Model

Archetypical example for classical magnetic systems

$$H = -J \sum_{\langle ij \rangle} s_i s_j$$

$s_i = \pm 1$ Ising spins.

$\langle ij \rangle$ sum over nearest-neighbours on the lattice.

$J > 0$ ferromagnetic coupling constant.

critical temperature $T_c > 0$ for $d > 1$.

Evolution, coupling to bath

Monte Carlo rule $s_i \rightarrow -s_i$ accepted with

| | | |
|---------------------------|----|--------------------------|
| $p = 1$ | if | $\Delta E < 0$ |
| $p = e^{-\beta \Delta E}$ | if | $\Delta E > 0$ |
| $p = 1/2$ | if | $\Delta \mathcal{E} = 0$ |

Non-conserved order parameter dynamics [$\uparrow\downarrow$ towards $\uparrow\uparrow$] etc. allowed.

[$m = 0$ to $m = 2$]

Kinetic n -vector models

More general ferromagnetic models

$$H = -J \sum_{\langle ij \rangle} \mathbf{s}_i \cdot \mathbf{s}_j$$

$$J > 0$$

Ferromagnetic coupling

$$\sum_{\langle ij \rangle}$$

Sum over nearest-neighbours on a d -dim. lattice.

$$s_i = \pm 1$$

Ising spins.

$$\mathbf{s}_i = (s_i^x, s_i^y)$$

e.g., xy two-component spins.

$$\ell^d \phi(\mathbf{r}) = \sum_{i \in V_{\mathbf{r}}} \mathbf{s}_i$$

Coarse-grained field over the volume $V = \ell^d$

$$L$$

Linear size of the system $L \gg \ell \gg a$

$$T_c > 0$$

for $d > 1$ and $L \rightarrow \infty$.

Coupling to the bath mimicked by **Monte Carlo updates**

The quench

Change in conditions

- Choose initial conditions, e.g. from equilibrium at a very high temperature

disordered state

- Evolve, in contact with the bath at a

critical or lower critical temperature

- Let the dynamics take the system towards

equilibrium

under the new conditions.

Stochastic dynamics

Open systems: discrete & continuous

- **Microscopic**: identify the ‘smallest’ relevant variables in the problem (e.g., the spins) and propose stochastic updates for them (as the **Monte Carlo** or **Glauber** rules for **spins**)
- **Coarse-grained**: write down a stochastic differential equation for the field, $\phi(\mathbf{r}, t)$, such as the **effective (Markov) Langevin equation**

$$\underbrace{m\ddot{\phi}(\mathbf{r}, t)}_{\text{Inertia}} + \underbrace{\gamma_0\dot{\phi}(\mathbf{r}, t)}_{\text{Dissipation}} = \underbrace{F(\phi)}_{\text{Deterministic}} + \underbrace{\xi(\mathbf{r}, t)}_{\text{Noise}}$$

with $F(\phi) = -\delta\mathcal{F}(\phi)/\delta\phi$ (with \mathcal{F} with a double well potential)


e.g., time-dependent stochastic scalar Ginzburg-Landau equation or the stochastic Gross-Pitaevskii equation

Ginzburg-Landau

Continuous scalar statistical field theory

Coarse-grain the spin

$$\phi(\mathbf{r}) = V_{\mathbf{r}}^{-1} \sum_{i \in V_{\mathbf{r}}} s_i.$$



The partition function is $\mathcal{Z} = \int \mathcal{D}\phi e^{-\beta \mathcal{F}(\phi)}$ with

$$\mathcal{F}(\phi) = \int d^d r \left\{ \frac{1}{2} [\nabla \phi(\mathbf{r})]^2 + \frac{T-J}{2} \phi^2(\mathbf{r}) + \frac{g}{4} \phi^4(\mathbf{r}) \right\}$$

Elastic + potential energy with the latter inspired by the results for the fully-connected model (entropy around $\phi \sim 0$ and symmetry arguments)


Uniform saddle point in the $V \rightarrow \infty$ limit : $\phi_{sp}(\mathbf{r}) = \langle \phi(\mathbf{r}) \rangle = \phi_0$

The free-energy density is $\lim_{V \rightarrow \infty} f_V(\beta, J, g) = \lim_{V \rightarrow \infty} V^{-1} \mathcal{F}(\phi_0)$


2nd order phase-transition

Bi-valued equilibrium states related by symmetry

\mathcal{F}



Ginzburg-Landau free-energy



Scalar order parameter

e.g. Ising magnets

Models

Summary : discrete vs. continuous

Ising spin models

$$H = -J \sum_{\langle ij \rangle} s_i s_j$$

$$\text{NCOP} \quad [\uparrow\downarrow \mapsto \uparrow\uparrow]$$

$$\text{COP} \quad [\uparrow\downarrow \mapsto \downarrow\uparrow]$$

Scalar field theories

$$\mathcal{F}[\phi] = \int d^d r \left[\frac{1}{2} (\nabla \phi)^2 - \frac{\mu}{2} \phi^2 + \frac{g}{4} \phi^4 \right]$$

$$\partial_t \phi(\mathbf{r}, t) = -\delta_{\phi(\mathbf{r}, t)} \mathcal{F}[\phi] + \xi(\mathbf{r}, t)$$

$$\partial_t \phi(\mathbf{r}, t) = -\nabla^2 \delta_{\phi(\mathbf{r}, t)} \mathcal{F}[\phi] + \eta(\mathbf{r}, t)$$

Overdamped limit is fine ; rescaling of time to eliminate γ_0

In the COP case $\langle \eta(\mathbf{x}, t) \eta(\mathbf{y}, t') \rangle = 2k_B T \nabla^2 \delta(\mathbf{x} - \mathbf{y}) \delta(t - t')$


Generalisations for vector cases. **Quenched disorder** can be introduced by taking the J_{ij} or the parameters in the field theory, e.g. μ , from a pdf.

Plan of this lecture


1. The phenomenon
2. Theoretical setting
3. **Critical and sub-critical quenches**
4. Dynamic scaling
5. Dynamic universality classes
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. Mean-field models
9. Modern studies

The problem


Up & down spins in a $2d$ Ising model: Instantaneous quench



$$T \rightarrow \infty$$



$$T = T_c$$




$$T < T_c$$

Question : starting from equilibrium at $T_0 \rightarrow \infty$ or $T_0 = T_c$ how is equilibrium at $T = T_c$ or $T < T_c$ attained ?

Real space viewpoint

$2d$ Ising model

Snapshots after an instantaneous quench from $T \rightarrow \infty$ to $T < T_c$



At $T = T_c$ **critical dynamics**

At $T < T_c$ **coarsening**

A certain number of **interfaces** or **domain walls** in the last snapshots.

Domain growth

- At $T = T_c$ the system needs to grow structures of all sizes.

Critical dynamics.

- At $T < T_c$: the system tries to order locally in one of the two competing equilibrium states at the new conditions.

Sub-critical coarsening.

The size of the equilibrated patches increases in time

- The relaxation time t_r needed to reach $\langle \phi \rangle_{eq}(T/J)$ diverges with the size of the system, $t_r(T/J, L) \rightarrow \infty$ when $L \rightarrow \infty$ for $T \leq T_c$
- **Dissipative dynamics** $dE/dt < 0$

Statement

In both cases one sees the growth of ‘red and white’ patches and **interfaces** surrounding such geometric domains.

Spatial regions of local equilibrium (with vanishing, at T_c , or non-vanishing, at $T < T_c$, order parameter) grow in time and

a single **growing length** $\mathcal{R}(t, T)$ can be identified and will be at the heart of dynamic scaling.



Here and in the following we measure T in units of J

Plan of this lecture

1. The phenomenon
2. Theoretical setting
3. Critical and sub-critical quenches
4. **Dynamic scaling**
5. Dynamic universality classes
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. Mean-field models
9. Modern studies

Critical dynamics



$$C(r, t) \equiv \langle s_i(t) s_j(t) \rangle_{|\mathbf{r}_i - \mathbf{r}_j| = r}$$



- Black curve : equilibrium relaxation, $r^{2-d-\eta}$.
- Coloured curves are for different times after the quench and they slowly approach the equilibrium one.
- From, say, $C(\mathcal{R}_c(t), t) = 1/e$ one gets $\mathcal{R}_c(t) \simeq t^{1/z_c}$ with z_c a critical exponent. (Other prescriptions give equivalent results.)
- z_c from t -dependent RG, $\simeq 2.17$ 2dIM Janssen, Schaub, Schmittmann 89

Sub-critical dynamcis

$$C(r, t) \equiv \langle s_i(t) s_j(t) \rangle_{|\mathbf{r}_i - \mathbf{r}_j| = r}$$



$$\langle \phi(t) \rangle = 0 \quad C_{eq}^c(r) \simeq e^{-r/\xi_{eq}}$$

- Coloured curves are $C(r, t)$ for different times after the quench.
- The growing length is $\mathcal{R}(t, T) \simeq t^{1/z_d}$ with $z_d = 2$
- $\mathcal{R}(t, T)$ is the averaged linear size of the domains.

Dynamic scaling

To be completed later

At late times there is a single *length-scale*, the *typical radius of the equilibrium structures (domains below T_c)* $\mathcal{R}(t, T)$, such that the structure is (in statistical sense) independent of time when lengths are scaled by $\mathcal{R}(t, T)$, e.g.

$$C(r, t) \equiv \langle s_i(t) s_j(t) \rangle |_{|\mathbf{x}_i - \mathbf{x}_j| = r} \sim f \left(\frac{r}{\mathcal{R}(t, T)} \right),$$

$$C(t, t_w) \equiv \langle s_i(t) s_i(t_w) \rangle \sim f_C \left(\frac{\mathcal{R}(t, T)}{\mathcal{R}(t_w, T)} \right),$$

etc. when $L \gg \mathcal{R} \gg \xi(T)$, $t, t_w \gg t_0$ and C small enough.


Suggested by experiments and numerical simulations **Lebowitz et al** 70s

Proved for a few cases.

Review Bray 94

Dynamic scaling

in phase ordering kinetics



Growing length $\mathcal{R}(t)$ and equilibrium reached for $\mathcal{R}(t_{eq}) \simeq L$

Typically $\mathcal{R}(t) \simeq t^{1/z}$ and $t_{eq} \simeq L^z$

Excess energy w.r.t. the equilibrium one stored in the domain walls ; $\Delta\mathcal{E}(t) \simeq \mathcal{R}^{-a}(t)$


Dynamic scaling

Quench of the $2d$ IM with NCOP from $T_0 \rightarrow \infty$ to $T < T_c$

Scaling regime $a \ll r \ll L$, $r \simeq \mathcal{R}(t, T) \simeq t^{1/z_d}$

$$C(r, t) \simeq m_{eq}^2(T) f_c \left(\frac{r}{\mathcal{R}(t, T)} \right)$$

Scaling looks perfect



$r/\mathcal{R}(t, T)$

Critical vs sub-critical

Space-time correlation: separation of time-scales & scaling

Critical quench

$$C(r, t) \simeq C_{eq}(r) f\left(\frac{r}{\mathcal{R}_c(t)}\right)$$

$$C_{eq}(r) \simeq r^{2-d-\eta}, \lim_{x \rightarrow 0} f(x) = 1 \text{ and } \lim_{x \rightarrow \infty} f(x) = 0.$$

Sub-critical

$$C(r, t) \simeq [C_{eq}(r) - m_{eq}^2] + m_{eq}^2 f\left(\frac{r}{\mathcal{R}(t, T)}\right)$$

$$C(0, t) = 1 \quad \forall t, \lim_{r \rightarrow 0} C_{eq}(r) = 1, \lim_{r \rightarrow \infty} C_{eq}(r) \propto \langle s_i \rangle_{eq}^2 = m_{eq}^2, \\ \lim_{x \rightarrow 0} f(x) = 1 \text{ (long times) and } \lim_{x \rightarrow \infty} f(x) = 0 \text{ (long distances).}$$

Plan of this lecture

1. The phenomenon
2. Theoretical setting
3. Critical and sub-critical quenches
4. Dynamic scaling
5. **Dynamic universality classes**
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. Mean-field models
9. Modern studies

Growing length

Dynamic universality classes

Use z_c in the growing length $\mathcal{R}_c(t)$ to identify critical dynamics universality classes.

But also, use the growing length $\mathcal{R}(t, T)$ to identify coarsening dynamic universality classes.

They depend on the dimension of the order parameter and the dynamic mechanism of growth (intimately related to the conservation laws).

Growing length

Dynamic universality classes at the critical point

At T_c , dynamic RG techniques work very well.

U. C. Tauber, *Critical Dynamics : A Field Theory Approach to Equilibrium and Non-Equilibrium Scaling Behavior* (Cambridge University Press, 2014)

One finds dynamic scaling, with the growing length

$$\mathcal{R}_c(t) \simeq t^{1/z_c}$$

z_c can be computed with methods that are very similar to critical exponents in static phase transitions.

Dynamic universality classes classified by the z_c values.

The scaling functions can be estimated as well

Growing length

Dynamic universality classes below the critical point

No systematic method

Focus on the dynamic mechanisms

Curvature driven

Numerical solution of the time-dependent Ginzburg-Landau equation




A. Langins (1st year master project)

$$\mathcal{R}(t, T) \simeq \lambda(T)t^{1/2}$$

Allen & Cahn, late 70s


Scalar field w/NCOP dynamics

- Curvature driven ($T = 0$): $\mathbf{v} \equiv \frac{d\mathbf{n}}{dt} \propto K \hat{\mathbf{n}}$ with $K = \nabla \cdot \hat{\mathbf{n}}$



Allen & Cahn 79



- Domain wall roughening ($T > 0$)
- Domain wall roughening and pinning by quenched disorder



e.g. elastic line in random media. **Kolton et al 05**

Curvature driven

Experiments with chiral liquid crystals



Scaling variable $r/\mathcal{R}(t) = r/t^{1/2}$


- 90°-Twisted in a confined space

Effectively 2D system

Many degrees of freedom

R. de Almeida PhD 19 Tokyo Univ.

R. de Almeida & K. A. Takeuchi 19



Curvature driven

Experiments with chiral liquid crystals

- Define a time series: $1/z_{\text{eff}}(t) \equiv \frac{d[\ln \rho(t)]}{d[\ln(t)]}$


- Average over $t \in [2.66\text{s} - 198.0\text{s}]$:

$$1/z = 0.52(3)$$

- Prior assumption of a power law:

$$1/z = 0.509(1)$$


$$t \in [2.66\text{s} - 198.0\text{s}]$$



- 90°-Twisted in a confined space

Effectively 2D system

Many degrees of freedom




R. de Almeida PhD 19 Tokyo Univ.

R. de Almeida & K. A. Takeuchi 19


Phase separation

Demixing transitions

Two species ● and ●, repulsive interactions between them.



Sketch



Experimental phase diagram

Binary alloy, **Hansen & Anderko, 54**


Phase separation

Spinodal decomposition in binary mixtures

A species \equiv spin up ; B species \equiv spin down

$2d$ Ising model with Kawasaki dynamics at T

locally conserved order parameter



50 : 50 composition Rounder boundaries

$$\mathcal{R}(t, T) \simeq \lambda(T) t^{1/3}$$

Huse 93

Weak disorder

e.g., random ferromagnets

At short time scales the dynamics is relatively fast and independent of the quenched disorder :

$$\mathcal{R}(t, T) \simeq \lambda(T) t^{1/z_d}$$

At longer time scales **domain-wall pinning** by disorder (J) dominates

Assume the wall has to overcome a length-dependent **barrier** scaling as

$B(\mathcal{R}) \simeq \Upsilon(T) \mathcal{R}^\psi$ to advance

The **Arrhenius time** needed to go over such a barrier is $t_A \simeq \tau_0 e^{\frac{B(\mathcal{R})}{k_B T}}$

This implies

$$\mathcal{R}(t, T) \simeq \left(\frac{k_B T}{\Upsilon(T)} \ln \frac{t}{\tau_0} \right)^{1/\psi}$$



Planar magnets

Schrielen pattern : gray scale according to $\sin^2 2\theta_i(t)$

Spin-waves



Vortices




After a quench vortices
annihilate & bind in pairs

$$\mathcal{R}(t, T) \simeq \lambda(T) [t / \ln(t/t_0(T))]^{1/2}$$

Frustrated magnets

e.g., $2d$ spin ice or vertex models





Stripe growth in the FM phase

Anisotropic growth, $\mathcal{R}_{\perp}(t, T)$ and $\mathcal{R}_{\parallel}(t, T)$

Polaritons

Dissipative & driven Gross-Pitaevskii equation



$$\mathcal{R}(t, T) \simeq \lambda(T) [t / \ln(t/t_0(T))]^{1/2}$$

Universality classes

as classified by the growing length

$$\mathcal{R}(t, T) \simeq \begin{cases} \lambda(T) t^{1/2} & \text{scalar NCOP} & z_d = 2 \\ \lambda(T) t^{1/3} & \text{scalar COP} & z_d = 3 \\ \lambda(T) \left(\frac{t}{\ln t} \right)^{1/2} & \text{planar NCOP in } d = 2 \\ \lambda(T) (\ln t)^{1/\psi} & \text{weak disorder NCOP} \end{cases}$$

Are scaling functions independent of
temperature, other parameters, microscopic dynamics ?

Super-universality ?

Dynamic scaling

Scaling functions

very early MC simulations **Lebowitz et al 70s** & experiments

One identifies a **growing linear size of equilibrated patches**

$$\mathcal{R}(t, T)$$

If this is the **only** length governing the dynamics, the **space-time correlation functions** should scale with $\mathcal{R}(t, T)$ according to

$$\text{At } T = T_c \quad C(r, t) \simeq C_{eq}(r) f_c\left(\frac{r}{\mathcal{R}_c(t)}\right)$$

Scaling fct f_c ✓

$$\text{At } T < T_c \quad C(r, t) \simeq C_{eq}^c(r) + m_{eq}^2 f\left(\frac{r}{\mathcal{R}(t, T)}\right)$$

Scaling fct f ?

Reviews **Hohenberg & Halperin 77** (critical) **Bray 94** (sub-critical)


Plan of this lecture

1. The phenomenon
2. Theoretical setting
3. Critical and sub-critical quenches
4. Dynamic scaling
5. Dynamic universality classes
6. **Two-time correlations and ageing**
7. Two-time responses and loss of memory
8. Mean-field models
9. Modern studies

Two-time self-correlation

e.g., MC simulation of the $2d$ IM at $T < T_c$

$$C(t, t_w) = N^{-1} \sum_{i=1}^N \langle s_i(t) s_i(t_w) \rangle$$



Stationary relaxation

Aging decay

Separation of time-scales : stationary – aging


$$C(t, t_w) = C_{st}(t - t_w) + m_{eq}^2 f \left(\frac{\mathcal{R}(t, T)}{\mathcal{R}(t_w, T)} \right)$$

$$C_{st}(0) = 1 - m_{eq}^2, \lim_{x \rightarrow \infty} C_{st}(x) = 0, f(1) = 1, \lim_{x \rightarrow \infty} f(x) = 0.$$


Two-time self-correlation

Comparison

Critical coarsening ($T = T_c$)



Sub-critical coarsening ($T < T_c$)



Separation of time-scales

Multiplicative

Additive

Ageing

Older samples relax more slowly

Older samples need more time to relax

spontaneously (correlation functions)





after a change in conditions (response functions)

t_w is the time that **measures the age of the system**

Huge literature on this phenomenology. Some reviews of experimental measurements were written by **Struick** on polymer glasses, **Vincent et al.** & **Nordblad et al.** on spin-glasses, **McKenna et al.** on all kinds of glasses.

Two-time self-correlation

Comparison




Thiospinel (spin-glass)

Lennard-Jones mixture (glass)

Plan of this lecture

1. The phenomenon
2. Theoretical setting
3. Critical and sub-critical quenches
4. Dynamic scaling
5. Dynamic universality classes
6. Two-time correlations and ageing
7. **Two-time responses and loss of memory**
8. Mean-field models
9. Modern studies

Response to perturbations



The **perturbation** couples **linearly** to the observable $B[\{s_i\}]$

$$H \rightarrow H - hB[\{s_i\}]$$

The **linear instantaneous response** of another observable $A[\{s_i\}]$ is

$$R_{AB}(t, t_w) \equiv \left\langle \left. \frac{\delta A[\{s_i\}](t)}{\delta h(t_w)} \right|_{h=0} \right\rangle$$

The **linear integrated response** or **dc susceptibility** is

$$\chi_{AB}(t, t_w) \equiv \int_{t_w}^t dt' R_{AB}(t, t')$$

Linear response

Critical and sub-critical coarsening

Critical coarsening

$$\chi(t, t_w) = \beta - \chi_{eq}(t - t_w) g \left(\frac{\mathcal{R}(t, T)}{\mathcal{R}(t_w, T)} \right)$$

Sub-critical coarsening

$$\chi(t, t_w) = \chi_{eq}(t - t_w) + [\mathcal{R}(t_w, T)]^{-a_\chi} g \left(\frac{\mathcal{R}(t, T)}{\mathcal{R}(t_w, T)} \right)$$

In both cases : $\chi_{eq}(t - t_w) = -(k_B T)^{-1} dC_{eq}(t - t_w)/d(t - t_w)$.


To be proven in the 3rd Lecture

Reviews

Crisanti & Ritort 03, Calabrese & Gambassi 05, Corberi *et al.* 07, LFC 11


Linear response

Sub-critical coarsening in the MC dynamics of 2dIM




Linear response

Coarsening vs glassy



Lippiello, Corberi & Zannetti 05



Sketch Chamon & LFC 07

There is no (weak) long-term memory in the coarsening problem. Just the stationary part will remain asymptotically, contrary to the sketch on the right for glasses & spin-glasses.

Plan of this lecture

1. The phenomenon
2. Theoretical setting
3. Critical and sub-critical quenches
4. Dynamic scaling
5. Dynamic universality classes
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. **Mean-field models**
9. Modern studies

The spherical $p = 2$ model

$$H = - \sum_{ij} J_{ij} s_i s_j + z \left(\sum_i s_i^2 - N \right)$$


Fully connected interactions

Gaussian distributed

interaction strengths J_{ij}

Spherical spins $\sum_i s_i^2 = N$

z is a Lagrange multiplier



$$\rho(\lambda_\mu) \propto \sqrt{(2J)^2 - \lambda_\mu^2}$$

$$H = - \sum_{\mu} \lambda_{\mu} s_{\mu}^2 + z \left(\sum_{\mu} s_{\mu}^2 - N \right)$$

Key: the largest eigenvalue
becomes **diffusive**,
 $k \approx \lambda_{\max} - z_{\infty} = 0$

Same scaling laws for two-time corr. and resp. but no space dependence

The $O(N)$ model

Upgrade the field to a vector $\phi \mapsto \boldsymbol{\phi}$ with $a = 1, \dots, N$ components

$$\boldsymbol{\phi} = (\phi_1, \dots, \phi_N)$$

The (over-damped) Ginzburg-Landau equation is now

$$\gamma_0 \partial_t \phi_a(\mathbf{r}, t) = - \frac{\delta \mathcal{F}[\boldsymbol{\phi}]}{\delta \phi_a(\mathbf{r}, t)} + \xi_a(\mathbf{r}, t)$$

The $N \rightarrow \infty$ limit allows one to decouple the vector components :

$$\phi_a(\mathbf{r}, t) [\mu - \frac{1}{N} \sum_{b=1}^N \phi_b^2(\mathbf{r}, t)] \mapsto \phi_a(\mathbf{r}, t) z(t)$$

and the equations are now linear with a global constraint.

Coarsening is linked to the growth of the **diffusive** $k = 0$ mode.

The $O(N)$ model

Upgrade the field to a vector $\phi \mapsto \boldsymbol{\phi}$ with $a = 1, \dots, N$ components

$$\boldsymbol{\phi} = (\phi_1, \dots, \phi_N)$$

The equations are now linear with a global constraint

$$\gamma_0 \partial_t \phi_a(\mathbf{r}, t) = \nabla^2 \phi_a(\mathbf{r}, t) + z(t) \phi_a(\mathbf{r}, t) + \xi_a(\mathbf{r}, t)$$

and

$$z(t) = \mu - N^{-1} \sum_a \phi_a^2(\mathbf{r}, t)$$

Solve for $\phi_a(\mathbf{r}, t)$ as a function of $z(t)$ and then impose the constraint to fix $z(t)$.

Coarsening is linked to the growth of the **diffusive** $k = 0$ mode, i.e. tendency to homogeneous order.

Summary

- At and below T_c growth of equilibrium structures.

- The linear size of the equilibrium patches is measured by $\mathcal{R}(t, T)$

- At T_c vanishing order parameter

Multiplicative scaling

$$C \simeq C_{eq} C_{ag}; \chi \simeq \chi_{eq} \chi_{ag}$$

- Below T_c non-vanishing order parameter

Additive scaling

$$C \simeq C_{eq} + C_{ag}; \chi \simeq \chi_{eq} + \chi_{ag}$$

- In both cases C_{ag} is finite while χ_{ag} vanishes asymptotically.

We shall discuss χ and how it compares to C later.

Phase ordering kinetics

The lecture was about

- Growth of equilibrium patches at T_c and below T_c .
- Divergence of $t_{eq}(L)$ with the system size.
- Existence of a single growing length $\mathcal{R}(t, T)$
- Separation of time-scales and dynamic scaling, e.g. $C = C_{eq} + C_{ag}$.
- Two kinds of correlations : Space-time and two-time ones.
- Dynamic universality classes at and below T_c .
- The more tricky/rich linear susceptibility.

Is there a static growing length in all systems with slow dynamics ?

Which one ?


Plan of this lecture

1. The phenomenon
2. Theoretical setting
3. Critical and sub-critical quenches
4. Dynamic scaling
5. Dynamic universality classes
6. Two-time correlations and ageing
7. Two-time responses and loss of memory
8. Mean-field models
9. **Modern studies**

Multiplicative noise


Numerical integration of the scalar field equations

NCOP



$$\mathcal{R}(t, T) \simeq t^{1/2}$$

COP



$$\mathcal{R}(t, T) \simeq t^{1/3}$$

The Voter Model

Archetypical example of opinion dynamics

Similar questions can be asked in very well-known problems in math, *e.g.*

Dynamics of a voter model starting from a random initial condition

- Purely dynamic, violation of detailed balance, no phase transition
- Two absorbing states
- The **dynamic mechanism** towards absorption is understood
domain growth is driven by interfacial noise

The Voter Model

Archetypical example of opinion dynamics

H does not exist - kinetic model

$s_i = \pm 1$ Ising spins that

sit on the vertices of a lattice.

Voter update rule

choose a spin at random, say s_i

choose one of its $2d$ neighbours at random, say s_j


set $s_i = s_j$

In two dimensions full consensus, *i.e.* $m = L^{-d} \sum_{i=1}^{L^d} s_i = \pm 1$ is reached in a timescale $t_C \simeq L^2$ (with $\ln L$ corrections)

Clifford & Sudbury 73, Holley & Liggett 75, Cox & Griffeaths 86

Ising vs. Voter

$s_i = \pm 1$ at $t = 0$ MCs, snapshots at $t = 4, 64, 512, 4096$ MCs



Ising


$T = 0$

T_c

Voter


Percolation issues

2d square IM at $T=0$




$t=0.0$

2d square IM at $T=0$




$t=0.57533$

2d square IM at T=0




$t=0.94844$

2d square IM at T=0




$t=2.00847$

2d square IM at T=0




$t=2.57898$

2d square IM at T=0




$t=3.99211$

2d square IM at $T=0$



$t=6.58423$


2d square IM at $T=0$




$t=7.46144$

2d square IM at $T=0$

The percolating structure was decided at $t_p \simeq 8$ MCs



$t=7.46144$



$t=128.0$

Complex field & cold atoms

Complex field theory in $3d$

Relativistic bosons; ^4He , type II superconductors, cosmology, etc.

$$-c^{-2}\ddot{\psi} + \nabla^2\psi + 2i\mu\dot{\psi} = g(\psi^2 - \rho)\psi$$

c is the velocity of light, ρ and g parameters in (Mexican hat) potential.

Limits

$\mu \rightarrow 0 :$

$$-c^{-2}\ddot{\psi} + \nabla^2\psi = g(|\psi|^2 - \rho)\psi$$

Goldstone

$c \rightarrow \infty :$

$$2i\mu\dot{\psi} + \nabla^2\psi = g(|\psi|^2 - \rho)\psi$$

Gross-Pitaevskii

models

Complex field theory in $3d$

Relativistic bosons; ^4He , type II superconductors, cosmology, etc.

$$-c^{-2}\ddot{\psi} + \nabla^2\psi + 2i\mu\dot{\psi} = g(\psi^2 - \rho)\psi$$

The energy functional

$$E = \int d^3x \left(c^{-2}|\dot{\psi}|^2 + |\nabla\psi|^2 - g\rho\psi^2 + g\psi^4 \right)$$

is conserved under the dynamics.

The energy is minimised by the static configuration $\psi = \sqrt{\rho} e^{i\chi}$ with $\chi = ct$

There are static vortex solutions, e.g. $\psi(\mathbf{x}) = f(r) e^{in\theta}$ with $f(0) = 0$ and $f(r \rightarrow \infty) = \sqrt{\rho}$, and $n \in \mathbb{Z}$ (thin tubes at the centre of which the field vanishes and the phase turns around).

Tsubota, Kasamatsu & Kobayashi 13, Kobayashi & Nitta 15, etc.

Complex field theory in 3d

Stochastic noise and dissipation added

$$-c^{-2}\ddot{\psi} + \nabla^2\psi + 2i\mu\dot{\psi} - \gamma\dot{\psi} = g(\psi^2 - \rho)\psi - \sqrt{\gamma T}\xi$$

Langevin-like dynamics

$-\gamma$ viscosity, ξ complex Gaussian white noise in normal form

$$\langle \xi_i(\mathbf{x}, t) \rangle = 0 \text{ and } \langle \xi_i(\mathbf{x}, t_1) \xi_j(\mathbf{y}, t_2) \rangle = \delta_{ij} \delta^{(3)}(\mathbf{x} - \mathbf{y}) \delta(t_1 - t_2)$$

Passage to Fokker-Planck formalism allows to show that the dynamics takes the system to

$$\lim_{t \rightarrow \infty} P(\psi, t) = P_{GB}(\psi) \propto e^{-\beta E}$$

Complex field theory in $3d$

Relativistic bosons; ^4He , type II superconductors, cosmology, etc.

$$-c^{-2}\ddot{\psi} + \nabla^2\psi + 2i\mu\dot{\psi} - \gamma\dot{\psi} = g(\psi^2 - \rho)\psi - \sqrt{\gamma T}\xi$$

Langevin-like dynamics

$-\gamma$ viscosity, ξ Gaussian white noise in normal form

In the limit $c \rightarrow \infty$, the stochastic Gross-Pitaevskii equation

$$(2i\mu - \gamma)\dot{\psi} = -\nabla^2\psi + g(\psi^2 - \rho)\psi + \sqrt{\gamma T}\xi$$

Gardiner et al 00s

3d XY lattice model

Archetypical classical magnetic example

$$H = -J \sum_{\langle ij \rangle} \mathbf{s}_i \cdot \mathbf{s}_j$$

$J > 0$ ferromagnetic coupling constant.

$\langle ij \rangle$ sum over nearest-neighbours on a 3d lattice

\mathbf{s}_i planar spins: two components with constant modulus \Rightarrow angle θ_i .

Second order phase transition with spontaneous symm breaking at $T_c > 0$.


Order parameter: spin-alignment, $\mathbf{m} \equiv N^{-1} \sum_i \langle \mathbf{s}_i \rangle$.

No intrinsic spin dynamics, Monte Carlo rules mimic coupling to thermal bath.

Non-conserved order parameter dynamics [$\uparrow\downarrow$ towards $\uparrow\uparrow$] etc. allowed.

Statics

Phase transition and order parameter in the field equation



$$L^3 m = |\sum_{ijk} \langle \psi_{ijk} \rangle|$$

critical temperature

$$T_c = 2.26$$

critical exponent


$$\beta = 0.347$$

Kobayashi & LFC 16


T_c and critical exponents from kurtosis (Binder parameter), susceptibility, specific heat, etc. Values compatible w/results from simulations **Ballesteros et al. 96**, **Hasenbusch & Török 99** and ϵ expansion **Guida & Zinn-Justin 98**, **Täuber & Diehl 14** for models in the same universality class.

Vortex configurations


In equilibrium




$0.6 T_c$



$0.8 T_c$



T_c



$1.2 T_c$


Periodic boundary conditions (torus) implies that the vortex lines are closed, i.e. loops.

Stochastic reconnection rule.

All vortex loops in blue, the longest one in red.

Dynamics after a quench

with g the control parameter



In the picture: annealing with finite rate.


Infinitely fast quench: $T \gg T_c$ for $t < 0$ and $T = 0$ for $t > 0$

Complex field theory in $3d$


Progressive elimination of vortex loops after a quench

$T \gg T_c$


$T = 0$



$t = 0$



$t = 3$




$t = 5$

As $\rho_{\text{vortex}} \downarrow$ the reconnection rule loses importance

Slow cooling & Kibble-Zurek

Finite rate quenching protocol

How is the scaling modified for a very slow quenching rate?



$$\Delta g \equiv g(t) - g_c = -t/\tau_Q \quad \text{with} \quad \tau_{Q1} < \tau_{Q2} < \tau_{Q3} < \tau_{Q4}$$


Standard time parametrization

$$g(t) = g_c - t/\tau_Q$$

Simplicity argument: linear cooling could be thought of as an approximation of any cooling procedure close to g_c .

Theoretical motivation

Network of cosmic strings



They should affect the Cosmic Microwave Background, double quasars, etc.

Picture from M. Kunz's group (Université de Genève)

Topological defects

instantaneous configurations



Domain walls in the $2d$ IM



Vortices in the $3d$ xy model

One can give a precise mathematical definition but the visual one is enough

Density of topological defects

Kibble-Zurek mechanics for 2nd order phase transitions

The three basic assumptions

- Defects are **created** close to the critical point.
- Their density in the ordered phase is inherited from the value it takes when the system falls out of equilibrium on the **symmetric** side of the critical point. It is determined by

Critical scaling above T_c

- The **dynamics in the ordered phase** is so slow that it can be **neglected**.

and one claim

- results are **universal**.

that we critically revisited within ‘thermal’ phase transitions

Topological defects

after an instantaneous quench : dynamic scaling




$$\Delta n(t) \simeq [\mathcal{R}(t, T)]^{-d} \simeq [\lambda(T(t))]^{-d} t^{-d/z_d}$$

Remember the initial ($g \rightarrow \infty$) configuration: already there !


Dissipative vs closed

One dimensional $\lambda\phi^4$ theory

$$H = a \sum_x \left\{ \frac{1}{2} \Pi_x^2 + \frac{1}{2a^2} (\phi_{x+a} - 2\phi_x)^2 + \lambda (\phi_x^2 - \varphi^2)^2 \right\}$$



Habib & Lythe 00



Bastianello *et al* in preparation